

Rational Use of Drugs: Eccentric or Non-existent?

Abdul Kader Mohiuddin *

Alumni, Faculty of Pharmacy, Dhaka University

*Correspondence Author: Abdul Kader Mohiuddin, Alumni, Faculty of Pharmacy, Dhaka University.

Received Date: September 28, 2022 | Accepted Date: October 10, 2022 | Published Date: October 25, 2022

Citation: Abdul Kader Mohiuddin, (2022) Rational Use of Drugs: Eccentric or Non-existent?. *International Journal of Clinical Research and Reports*.1(1); DOI:10.31579/2835-785X/003

Copyright: © 2022 Abdul Kader Mohiuddin, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Proper use of medicine or taking medicine in correct order is essential to cure any disease. Even patients from developed nations have trouble staying on top of their drug compliance. When it comes to improper medicine use, there is an odd parallel between underdeveloped, emerging nations and the so-called developed world in the West. The key factor influencing whether patients stick to their treatment plan is their understanding and perception of the disease.

Keywords: right administration; national health service

Introduction

Proper use of medicine or taking medicine in correct order is essential to cure any disease. Medication adherence usually refers to whether patients take their medications as prescribed. According to the WHO, lack of adherence to treatment regimens leads to major problems among patients, mostly with chronic illnesses. "Right administration" depends on at least 5 right factors--right patient, right drug, right time, right dose and right route [1]. "Medicines won't work if you don't take it right"--This simple fact is not understood by most people in the world, as a result still more than half of the patients with chronic diseases in the developed world do not take their medicine correctly--says WHO [2].

Patients with chronic diseases may find it particularly difficult to adhere because they frequently need to take their medications for an extended period of time, sometimes for the remainder of their lives. There are several reasons why patients may find it challenging to adhere to treatment regimens, and CDC estimates that medication non-compliance leads to 30 to 50% of chronic disease treatment failures. Poor adherence may cause treatment outcomes to not be achieved, symptoms to worsen, and one's health to deteriorate [3].

In the UK, up to 50% of medicines are not taken as intended and 60% of publicly funded healthcare systems of the UK National Health Service (NHS) patients failed to receive the right treatment within 18 weeks [4-6]. According to a New Zealand-based Journal medication non-adherence alone accounts for at least 10% of hospitalizations, \$300 billion in annual medical costs, and more than 1,25,000 deaths in the US alone [7]. A recent Canadian study found that 30% of patients stop taking their medication before it is instructed, and one in four do not fill their prescription or take less than prescribed [8]. Medication non-adherence contributes more than half of the \$500 billion total avoidable costs attributed to suboptimal medicine use globally each year [9].

A strange similarity can be found in under-developed, developing countries and the so-called developed world in the West or the Middle-East when it comes to not taking medicine properly. More than half of the antibiotics worldwide are sold without a prescription and CDC stated, 30-50% of antibiotics prescribed in hospitals are inappropriate or unnecessary [10, 11]. A recent study published by The Lancet, funded by the Bill & Melinda Gates Foundation, Wellcome Trust, states that nearly 5 million deaths worldwide in 2019 were related to bacterial resistance, which is expected to double by 2050 [12].

Globally, NSAIDs are responsible for at least 650,000 hospitalizations and 165,000 deaths annually [13]. Overuse of this class of drugs can cause kidney injury, and their side effects can be 3 to 4 times higher in kidney compromised patients [14]. Many studies have reported widespread misuse of these drugs in Dengue, Chikungunya and Covid-19 patients. Especially in Dengue or Covid-19 patients, it is more important to maintain the hydration level of the body than to bring down the fever with the pain killers. In children, the use of excess Paracetamol syrup or suppositories may cause stomach irritation, which hampers digestion and led to vomiting and ended up with hospitalization. Most hospitalizations or ICU admissions among those patients could be prevented, with few exceptions, simply by preventing dehydration at home with saline and fruit juice or simply by drinking more water.

More or less 40% of Covid-19 patients report sleep disturbances--use of Benzodiazepines in Covid-19 patients increases the incidence of delirium, depresses the system in patients with compromised respiratory functions, and contraindicated with some anti-viral medications [15, 16].

Around 300 metric tons of morphine-type painkillers are used worldwide each year, less than 1% of which distributed to low-and-middle income countries, says the American Journal of Public Health [17]. So their misuse and related side effects are also retained by the developed world.

According to a WHO report, only half of patients in developed countries adhere to treatment guidelines for chronic diseases, which is much less in developing countries [18]. Several studies among diabetic patients in South Asian countries have shown that nearly half of patients do not adhere to their prescribed medication and are at risk of acute and long-term complications, resulting in increased hospitalization rates and medical costs [19, 20].

Three-quarters of elderly patients worldwide are unable to adhere to appropriate long-term treatment regimens—due to multiple physical complications and additional medication burden [21]. Elderly patients taking at least 5 medications are at increased risk of mild cognitive impairment, dementia, falls, frailty, disability, and mortality, while ADEs are estimated to be 5% to 28% of acute geriatric medical admissions [22, 23].

A recent study by the American Heart Association revealed that patients with high blood pressure do not follow treatment guidelines because of--(1) suboptimal dosing or prescribing the wrong medication (2) lack of insurance or lack of health care access and (3) patient failure to comply prescribed medication or other lifestyle guidelines [24].

“Medical costs are barriers to adherence to proper clinical guidelines for long-term diseases in poor countries”-- although discussed in many forums but forgetfulness, confusion about the duration required for medication use and mistrust about the overall efficacy of medication are among the reasons for non-adherence to diabetes management protocols in Middle Eastern countries [25].

Epilepsy is still seen in most countries of the world as an evil spirit --although two-thirds of patients can become seizure-free with adequate treatment, poor adherence to proper guidelines is a major problem for effective recovery [26].

Several identified reasons for non-adherence to treatment guidelines for chronic diseases:

1. Patient's socio-economic status: low health literacy, lack of family or social support network, unstable living or homelessness, financial insecurity

2. Treatment-related: complexity and duration of treatment procedures, frequent changes in medication regimen, lack of immediate results, real or perceived unpleasant side effects, interference with lifestyle

3. Health system-related: high treatment costs, limited health system for patient education and follow-up, doctor-patient relationship, patient trust in health care, long waits, lack of patient information materials

4. Patient-related: visual-hearing and cognitive impairment, mobility and dexterity, psychological and behavioral factors, perceived risk of disease susceptibility, superstitions and stigmatization by disease, etc. [27]

Finally, it can be said that patients' knowledge and perception of the disease is the main driving force in determining their adherence to the treatment regimen. Health care providers should explore providing more effective health-education to identify patients' attitudes toward disease, trust in medications, psychological stressors, and increase medication adherence.

Acknowledgement

I'm thankful to Khulud Alosaimi, Department of Pharmacology and Toxicology, Faculty of Medicine, Umm Al-Qura University, Makkah for his valuable time to audit my paper and for his thoughtful suggestions. I'm also grateful to seminar library of Faculty of Pharmacy, University of Dhaka and BANSDOC Library, Bangladesh for providing me books, journal and newsletters.

Abbreviations:

Adverse drug events (ADEs)

Non-steroidal anti-inflammatory drugs (NSAIDs)

The Centers for Disease Control and Prevention (CDC)

World Health Organization (WHO)

Financial Disclosure or Funding: N/A

Conflict of Interest: The author declares that he has no competing interests.

Informed Consent: N/A

References

1. Grissinger M. (2010) The Five Rights: A Destination Without a Map. *P T*. 35(10):542.
2. Brown MT, Bussell JK. (2011) Medication adherence: WHO cares? *Mayo Clin Proc*. 86(4):304-314.
3. Neiman AB, Ruppert T, Ho M, Garber L, Weidle PJ, Hong Y, George MG, Thorpe PG. (2018) CDC Grand Rounds: Improving medication adherence for chronic disease management - Innovations and opportunities. *Am J Transplant*. 18(2):514-517.
4. Barnett, N L. (2014) “Medication Adherence: Where Are We Now? A UK Perspective.” *European Journal of Hospital Pharmacy*, vol. 21, no. 3, 19 pp. 181–184.,
5. Campbell, Denis, and Pamela Duncan. (2022) “Record 6.8m People Waiting for Hospital Treatment in England.” *The Guardian*.
6. Andrews, Luke. (2022) “Revealed: 60% of NHS Patients Needing Routine Procedures Have Waited Longer than 18 Weeks at England's Worst-Performing Hospital Trust.” *Mail Online*, [patients-waited-18-weeks-treatment-worst-performing-hospital-trust.html](https://www.mailonline.com/news/health/nhs-patients-waited-18-weeks-treatment-worst-performing-hospital-trust.html).
7. Cutler RL, Torres-Robles A, Wiecek E, Drake B, Van der Linden N, Benrimoj SI, Garcia-Cardenas V. (2019) Pharmacist-led medication non-adherence intervention: reducing the economic burden placed on the Australian health care system. *Patient Prefer Adherence*. 13:853-862.
8. Bonsu KO, Young S, Lee T, Nguyen H, Chitsike RS. (2022) Adherence to Antithrombotic Therapy for Patients Attending a Multidisciplinary Thrombosis Service in Canada - A Cross-Sectional Survey. *Patient Prefer Adherence*. 16:1771-1780.
9. Furniss D, Barber N, Lyons I, Eliasson L, (2014) Blandford A. Unintentional non-adherence: can a spoon full of resilience help the medicine go down? *BMJ Qual Saf*. 23(2):95-98.
10. Bahta M, Tesfamariam S, Weldemariam DG, Yemane H, Tesfamariam EH, Alem T, Russom M. (2020) Dispensing of antibiotics without prescription and associated factors in drug retail outlets of Eritrea: A simulated client method. *PLoS One*. 24;15(1):e0228013.
11. “CDC's 6/18 Initiative: Improve Antibiotic Use.” Centers for Disease Control and Prevention, Centers for Disease Control and Prevention, 9 Mar. 2022,
12. Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet*. 2022 Feb 12;399(10325):629-655.
13. Kasciūškevičiūtė S, Gumbrevičius G, Vendzelytė A, Ščiupokas A, Petrikonis K, Kaduševičius E. (2018) Impact of the World Health Organization Pain Treatment Guidelines and the European Medicines Agency Safety Recommendations on Nonsteroidal Anti-Inflammatory Drug Use in Lithuania: An Observational Study. *Medicina (Kaunas)*. May 11;54(2):30.
14. Lucas GNC, Leitão ACC, Alencar RL, Xavier RMF, Daher EF, Silva Junior GBD. (2019) Pathophysiological aspects of nephropathy caused by non-steroidal anti-inflammatory drugs. *J Bras Nefrol*.;41(1):124-130.
15. Jahrami H, BaHammam AS, Bragazzi NL, Saif Z, Faris M, Vitiello MV. (2021) Sleep problems during the COVID-19

- pandemic by population: a systematic review and meta-analysis. *J Clin Sleep Med*. 17(2):299-313.
16. Ostuzzi G, Papola D, Gastaldon C, Schoretsanitis G, Bertolini F, Amaddeo F, Cuomo A, Emsley R, Fagiolini A, Imperadore G, Kishimoto T, Michencigh G, Nosé M, Purgato M, Dursun S, Stubbs B, Taylor D, Thornicroft G, Ward PB, Hiemke C, Correll CU, Barbui C. (2020) Safety of psychotropic medications in people with COVID-19: evidence review and practical recommendations. *BMC Med*. 18(1):215.
 17. Bhadelia A, De Lima L, (2019) Arreola-Ornelas H, Kwete XJ, Rodriguez NM, Knaul FM. Solving the Global Crisis in Access to Pain Relief: Lessons From Country Actions. *Am J Public Health*. 109(1):58-60.
 18. Chauke GD, Nakwafila O, Chibi B, Sartorius B, (2022) Mashamba-Thompson T. Factors influencing poor medication adherence amongst patients with chronic disease in low-and-middle-income countries: A systematic scoping review. *Heliyon*. 8(6):e09716.
 19. Chong E, Wang H, King-Shier KM, Quan H, Rabi DM, Khan NA. (2014) Prescribing patterns and adherence to medication among South-Asian, Chinese and white people with Type 2 diabetes mellitus: a population-based cohort study. *Diabet Med*.
 20. Sohal T, Sohal P, King-Shier KM, Khan NA. (2015) Barriers and Facilitators for Type-2 Diabetes Management in South Asians: A Systematic Review. *PLoS One*. 10(9):e0136202.
 21. Félix IB, Henriques A. (2021) Medication adherence and related determinants in older people with multimorbidity: a cross-sectional study. *Nurs Forum*. 56:834–843.
 22. Chippa, Venu, and Kamalika Roy.(2022) “Geriatric Cognitive Decline and Polypharmacy.” National Library of Medicine, StatPearls, Varghese D, Ishida C, Haseer Koya H. (2022) “Polypharmacy.” National Library of Medicine, StatPearls,
 23. Choudhry NK, Kronish IM, Vongpatanasin W, Ferdinand KC, Pavlik VN, Egan BM, Schoenthaler A,(2022) Houston Miller N, Hyman DJ; American Heart Association Council on Hypertension; Council on Cardiovascular and Stroke Nursing; and Council on Clinical Cardiology. Medication Adherence and Blood Pressure Control: A Scientific Statement From the American Heart Association. *Hypertension* 79(1):e1-e14.
 24. Alsairafi ZK, Taylor KM, Smith FJ, Alattar AT. (2016) Patients' management of type 2 diabetes in Middle Eastern countries: review of studies. *Patient Prefer Adherence*. 10:1051-1062.
 25. Lossius, M. I., Alfstad, K. Å., Aaberg, K. M., & Nakken, K. O. (2017). Seponering av Antiepileptika Ved anfallsfrihet – når og hvordan? *Tidsskrift for Den Norske Legeforening*, 137(6), 451–454.
 26. Jin J, Sklar GE, Min Sen Oh V, Chuen Li S. (2008) Factors affecting therapeutic compliance: A review from the patient's perspective. *Ther Clin Risk Manag*. 4(1):269-286.

Ready to submit your research? Choose ClinicSearch and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At ClinicSearch, research is always in progress.

Learn more <https://clinicsearchonline.org/journals/international-journal-of-clinical-research-and-reports>



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.