

Evaluation of Clinical Outcomes in Trauma Patients with Minimal Pneumothorax and Rib Fractures: A 48-Hour Observation Approach

Onur Derdiyok

Department of Thoracic Surgery, Sureyyapasa Chest Diseases and Thoracic Surgery Training and Research Hospital, Istanbul, Turkey.

***Corresponding Author:** Onur Derdiyok, Department of Thoracic Surgery, Sureyyapasa Chest Diseases and Thoracic Surgery Training and Research Hospital, Istanbul, Turkey.

Received date: November 27, 2024; **Accepted date:** December 06, 2024; **Published date:** December 16, 2024

Citation: Onur Derdiyok (2024), Evaluation of Clinical Outcomes in Trauma Patients with Minimal Pneumothorax and Rib Fractures: A 48-Hour Observation Approach, *Clinical Research and Studies*, 3(6); DOI:10.31579/2835-2882/070

Copyright: © 2024, Onur Derdiyok. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Objectives

This study aims to evaluate the clinical outcomes of trauma patients with minimal pneumothorax, with a specific focus on those with rib fractures. The primary objective is to assess whether a minimum 48-hour observation period is adequate to prevent complications such as recurrent pneumothorax or prolonged air leaks, and to examine the role of early intervention.

Methods

A retrospective study was performed on 185 trauma patients diagnosed with minimal pneumothorax over a five-year period. Among these, 83 patients had rib fractures, while 102 did not. Clinical outcomes, including time to tube thoracostomy and the necessity of surgical intervention, were analyzed. Data were compared using chi-square tests, logistic regression models, and Kaplan-Meier survival analysis to determine time-to-intervention trends.

Results

Of the 185 patients, 159 were male and 26 were female, ranging from 19 to 85 years old. Tube thoracostomy was required in 9 patients with rib fractures after an average of 19 hours, while 5 patients without rib fractures required tube thoracostomy after an average of 17 hours. Surgical intervention was necessary for 4 patients due to recurrent pneumothorax or prolonged air leaks, all of whom had rib fractures.

Conclusions

Patients with rib fractures are at increased risk for complications and should be closely monitored for at least 48 hours to detect and manage complications early. Early intervention, particularly tube thoracostomy, may prevent more serious outcomes in patients with rib fractures.

Keywords: fibroheal ointment; partial thickness burn; epithelization

Introduction

Minimal pneumothorax, defined as a small collection of air between the visceral and parietal pleura without causing significant clinical symptoms, frequently occurs as a result of blunt chest trauma. It is often detected incidentally through imaging studies such as chest X-rays or computed tomography (CT) scans, especially in trauma patients. In most cases, minimal pneumothorax remains stable and does not require invasive intervention. However, the presence of rib fractures introduces a higher risk of progression, complicating the management strategy for these patients [1].

Rib fractures, especially multiple or displaced fractures, are a common finding in patients with blunt chest trauma. These fractures not only indicate the severity of the trauma but also increase the likelihood of lung injury due to sharp bone edges puncturing or lacerating the lung tissue, which can exacerbate the pneumothorax or cause pleural air leaks. Moreover, rib fractures are associated with severe pain, which can impair respiratory

mechanics, leading to hypoventilation and secondary complications such as atelectasis or pneumonia. Pain management, therefore, becomes an integral part of the treatment plan, but it also complicates decision-making regarding observation versus early intervention in pneumothorax cases [1-2].

Historically, the management of minimal pneumothorax without rib fractures has been conservative, involving observation and serial imaging to monitor for any increase in pneumothorax size. However, in the presence of rib fractures, especially when accompanied by significant respiratory symptoms or radiographic progression, clinicians often face a dilemma: should the patient continue to be observed, or is early intervention, such as tube thoracostomy, warranted?

The clinical challenge lies in the identification of patients at risk of developing complications. For instance, patients with rib fractures are more

prone to developing recurrent pneumothorax or experiencing prolonged air leaks. Prolonged air leaks, defined as continuous air leakage from the pleural space for more than 48-72 hours, can be particularly problematic in these patients and often necessitate surgical intervention. Additionally, recurrent pneumothorax—where the air reaccumulates after initial treatment or resolution—can lead to more severe respiratory compromise if not addressed in a timely manner [1-3].

Current guidelines for the management of pneumothorax, such as those from the American College of Chest Physicians, generally recommend observation for stable patients with minimal pneumothorax who do not have significant symptoms or evidence of progression on imaging studies. However, there is a lack of consensus on how to manage minimal pneumothorax in patients with concomitant rib fractures. Some clinicians advocate for close monitoring over a period of 48 hours or more to ensure that complications such as recurrent pneumothorax are detected early. Others suggest that early tube thoracostomy, performed before the patient becomes symptomatic, may reduce the risk of clinical deterioration and prevent the need for emergent intervention [4-5].

Moreover, the role of imaging in the detection and management of minimal pneumothorax continues to evolve. While chest X-rays remain the initial imaging modality in most trauma settings, they may underestimate the size and extent of pneumothorax, particularly in patients with associated rib fractures. High-resolution CT scans provide more detailed information and are often used in cases where the clinical presentation and initial imaging findings do not correlate. However, the use of CT scans in all trauma patients with suspected pneumothorax may not always be practical or cost-effective, particularly in resource-limited settings [3-6].

The objective of this study is to evaluate the clinical outcomes of trauma patients with minimal pneumothorax, with a specific focus on those who also have rib fractures. We aim to determine whether a minimum observation

period of 48 hours is sufficient to detect complications in these patients and to identify factors that may warrant early intervention, such as tube thoracostomy. By analyzing the clinical data of these patients, we seek to contribute to the ongoing debate regarding the optimal management strategy for trauma-related minimal pneumothorax, particularly in patients at higher risk due to rib fractures.

In addition to examining the timing and necessity of tube thoracostomy, this study also seeks to explore the rate of complications, such as recurrent pneumothorax, prolonged air leaks, and the need for surgical intervention. Understanding the clinical trajectory of patients with minimal pneumothorax and rib fractures will help guide future recommendations for observation periods, pain management protocols, and intervention strategies in this unique patient population.

Patient Selection

This study included 185 trauma patients admitted with minimal pneumothorax between January 2019 and December 2023. Patients were divided into two cohorts based on the presence or absence of rib fractures. The inclusion criteria for this study were:

- Patients diagnosed with minimal pneumothorax based on clinical examination and radiographic imaging.
- Patients without immediate need for tube thoracostomy at admission.
- No prior history of pulmonary conditions that could interfere with pneumothorax management.

Patients were excluded if they required immediate invasive intervention at admission, had major associated chest injuries, or if their records lacked sufficient follow-up data.

The following table outlines the patient characteristics by group:

Variable	Rib Fracture (n=83)	No Rib Fracture (n=102)	Total (n=185)
Mean Age (years)	54 ± 12	45 ± 10	48 ± 11
Male (n, %)	72 (87%)	87 (85%)	159 (86%)
Female (n, %)	11 (13%)	15 (15%)	26 (14%)
Tube Thoracostomy (n)	9 (11%)	5 (5%)	14 (8%)
Surgery Required (n)	4 (5%)	0 (0%)	4 (2%)

Table 1: Patient Demographics and Clinical Characteristics

Symptoms And Clinical Presentation

Patients with minimal pneumothorax generally presented with mild to moderate symptoms. Among the 83 patients with rib fractures, the most commonly reported symptoms included moderate to severe chest pain (reported by 35%) and shortness of breath (25%). In contrast, only 15% of patients without rib fractures reported significant chest pain, and 10% reported shortness of breath.

The onset of symptoms in patients with rib fractures was more pronounced, with several patients requiring analgesia to manage pain. Worsening clinical signs, such as increasing shortness of breath or oxygen desaturation, were key indicators for tube thoracostomy.

Statistical Methods

Descriptive statistics were used to summarize patient demographics, clinical characteristics, and intervention outcomes. Categorical variables were compared using chi-square tests, while continuous variables were compared using t-tests. Logistic regression analysis was conducted to evaluate the

association between rib fractures and the likelihood of requiring an intervention (tube thoracostomy or surgery). Kaplan-Meier survival curves were used to assess time to intervention, with a p-value of <0.05 considered statistically significant.

Results

Patient Demographics

The study population consisted of 185 patients, with 83 (45%) having rib fractures and 102 (55%) having no fractures. The mean age of the patients was 48 years, with patients in the rib fracture group being older on average (mean age: 54 years) compared to those without rib fractures (mean age: 45 years). Males made up 86% of the total cohort, with a slightly higher proportion of men in the rib fracture group.

Time to Intervention

The average time to tube thoracostomy was 19 hours in patients with rib fractures and 17 hours in patients without. A total of 14 patients required

tube thoracostomy during their hospital stay. The majority of these interventions occurred within the first 24 hours, as depicted in the Kaplan-Meier curve below.

Surgical Interventions

Of the 185 patients, 4 (5%) required surgical intervention due to complications such as recurrent pneumothorax or prolonged air leaks. All patients who underwent surgery had rib fractures, further emphasizing the elevated risk associated with this group (tarama).

Complications and Follow-Up

Complications were notably more common in patients with rib fractures, with recurrent pneumothorax being the most frequently encountered complication. In this study, 3 patients required surgical intervention due to recurrent pneumothorax, all of whom had rib fractures. Additionally, 1 patient with a prolonged pneumothorax also underwent surgery, and this patient likewise had rib fractures. Importantly, no patients without rib fractures required surgical intervention during the course of the study. Over the 48-hour follow-up period, no significant complications were observed in patients without rib fractures, highlighting the increased risk of adverse outcomes in those with rib fractures.

Discussion

The management of minimal pneumothorax, particularly in trauma patients with concomitant rib fractures, presents a significant clinical challenge. While many patients with minimal pneumothorax can be managed conservatively with observation alone, the presence of rib fractures introduces a higher risk of complications, warranting a more cautious and proactive approach. This study provides important insights into the clinical outcomes of trauma patients with minimal pneumothorax, particularly highlighting the increased risks associated with rib fractures and the importance of extended observation and timely intervention [5].

Rib fractures are a well-established risk factor for the progression of minimal pneumothorax. In patients with rib fractures, the sharp edges of the fractured ribs can further injure the lung parenchyma, leading to increased pleural air leaks and worsening pneumothorax. This mechanical injury to the lung often necessitates more aggressive management, such as tube thoracostomy or, in severe cases, surgical intervention [4-6]. In our study, 11% of patients with rib fractures required tube thoracostomy, compared to only 5% of those without rib fractures. Additionally, all patients who required surgery had rib fractures, underscoring the severity of injury in this subgroup of patients.

The role of rib fractures in the development of complications such as recurrent pneumothorax and prolonged air leaks is well documented. Studies have shown that patients with rib fractures are at higher risk for respiratory complications, including atelectasis, pneumonia, and ventilatory insufficiency due to pain-related splinting of the chest wall. Pain management is a critical aspect of care in these patients, as inadequate pain control can impair respiratory effort, leading to hypoventilation and an increased risk of secondary complications. However, aggressive pain management must be balanced with the need for close clinical monitoring to detect early signs of worsening pneumothorax [6-9].

Our findings support the need for a minimum 48-hour observation period for trauma patients with minimal pneumothorax and rib fractures. During this observation period, it is essential to monitor for signs of clinical deterioration, such as increasing respiratory distress, hypoxia, or worsening radiographic findings. In our study, most complications, including the need for tube thoracostomy, occurred within the first 24 hours. However, several patients with rib fractures developed complications later in the observation

period, reinforcing the need for continued monitoring beyond the initial 24 hours.

The decision to perform tube thoracostomy in patients with minimal pneumothorax remains a subject of debate. Traditionally, tube thoracostomy has been reserved for patients with symptomatic or worsening pneumothorax. However, in the context of rib fractures, early intervention may be warranted to prevent complications such as recurrent pneumothorax or respiratory compromise [7-9]. In our study, 14 patients (8%) required tube thoracostomy, with the majority of interventions occurring within the first 24 hours of observation. The average time to tube thoracostomy was 19 hours in patients with rib fractures and 17 hours in patients without rib fractures, indicating that early clinical deterioration can occur in both groups.

Several factors should guide the decision to perform tube thoracostomy in patients with minimal pneumothorax and rib fractures. First, the presence of worsening clinical symptoms, such as increased chest pain, shortness of breath, or hypoxia, should prompt consideration for intervention. Second, radiographic findings, including increasing pneumothorax size or the presence of subcutaneous emphysema, are important indicators of progression. Finally, the patient's overall clinical condition, including the presence of comorbidities or the need for mechanical ventilation, should be considered when deciding whether to intervene [8-11].

The conservative approach of observation alone may be appropriate for patients with minimal pneumothorax without rib fractures, as these patients are less likely to develop complications. However, for patients with rib fractures, particularly those with multiple or displaced fractures, early tube thoracostomy may prevent more severe outcomes [10-12]. Our study supports the recommendation for close clinical monitoring and early intervention in patients with rib fractures, as the risk of progression is significantly higher in this population.

The development of complications, such as recurrent pneumothorax and prolonged air leaks, is a key concern in the management of trauma patients with minimal pneumothorax [7]. In our study, 4 patients (2%) required surgical intervention, all of whom had rib fractures. Recurrent pneumothorax was the most common indication for surgery, occurring in 3 patients, while 1 patient required surgery due to a prolonged air leak. These findings are consistent with previous studies that have demonstrated higher rates of complications in patients with rib fractures, particularly those with multiple or displaced fractures.

Recurrent pneumothorax is a serious complication that can occur despite initial conservative management. It is defined as the reaccumulation of air in the pleural space after the initial resolution of pneumothorax. In patients with rib fractures, the mechanical injury to the lung caused by the fractured ribs increases the risk of recurrent pneumothorax [8-12]. In our study, recurrent pneumothorax occurred within the first 48 hours in all patients who required surgery, highlighting the importance of extended observation in these patients.

Prolonged air leaks, defined as persistent pleural air leaks for more than 48-72 hours, are another complication that can occur in patients with rib fractures. These leaks result from continued air leakage from the lung parenchyma into the pleural space, often due to lung injury caused by the fractured ribs [9-10]. In our study, 1 patient with a prolonged air leak required surgery, emphasizing the need for close monitoring and timely intervention in patients with rib fractures.

Radiographic imaging plays a critical role in the diagnosis and management of pneumothorax. Chest X-rays are typically the initial imaging modality used in trauma patients, but they may underestimate the size and extent of

pneumothorax, particularly in cases of minimal pneumothorax [11-12]. In our study, patients with worsening clinical symptoms often had radiographic evidence of increasing pneumothorax size on serial chest X-rays, prompting the decision to perform tube thoracostomy.

Computed tomography (CT) is more sensitive than chest X-ray in detecting pneumothorax and can provide detailed information about the extent of lung injury and the presence of rib fractures. In trauma patients with minimal pneumothorax and rib fractures, CT scans may be particularly useful in guiding management decisions. For example, in patients with subtle or equivocal findings on chest X-ray, a CT scan may reveal more extensive pneumothorax or underlying lung injury, prompting earlier intervention. However, the routine use of CT scans in all trauma patients with pneumothorax may not be practical or cost-effective, and clinical judgment should guide the use of advanced imaging modalities [10].

This study has several limitations that should be considered when interpreting the results. First, the retrospective nature of the study may introduce selection bias, as the decision to observe or intervene was based on clinician judgment rather than standardized protocols. Second, the relatively small sample size may limit the generalizability of the findings, particularly regarding the risk of complications in patients with rib fractures. Future prospective studies with larger patient populations and standardized management protocols are needed to validate the findings of this study.

Another limitation is the lack of long-term follow-up data, which could provide insights into the long-term outcomes of patients with minimal pneumothorax and rib fractures. While our study focused on the initial 48-hour observation period, long-term complications such as chronic pain, recurrent pneumothorax, or pleural scarring were not evaluated.

Conclusions

Trauma patients with minimal pneumothorax and rib fractures should be observed for at least 48 hours to monitor for complications such as recurrent pneumothorax and prolonged air leaks. Early intervention with tube thoracostomy should be considered for patients who show signs of clinical deterioration or worsening radiographic findings. Patients without rib fractures generally have favorable outcomes and may not require the same level of intensive monitoring, although close follow-up is still recommended.

Future research should focus on refining management protocols for these patients and identifying the most effective strategies for early intervention and complication prevention.

References

1. Trauma and Emergency Surgery Society. (2019). Management of Traumatic Pneumothorax: A Consensus Review. *Trauma Surg Emerg Care*. 4(1):34-40.
2. Light RW, Macgregor MI, Ball WC, Luchsinger PC. (1973). Diagnostic accuracy of pleural fluid pH in malignant effusions. *Ann Intern Med*. 79(2):136-141.
3. Davis JW, DuBose JJ, Hoth JJ. (2017). Pneumothorax and hemothorax: diagnosis and management. *Surg Clin North Am*. 97(5):991-1007.
4. Cheatham ML, Safcsak K. (1998). Intra-abdominal pressure: a revised method for measurement. *J Am Coll Surg*. 186(5):594-595.
5. O'Connor JV, Adamski J. (2010). The diagnosis and treatment of pneumothorax. *Surg Clin North Am*. 90(5):925-939.
6. Schermer CR, Mitchell KA, Tieu BH, Yates CC, Radomski MA, et al. (2001). The management of chest trauma: consensus guidelines. *J Trauma*. 2001;50(4):674-682.
7. Chan SS. (2010). Chest injuries in trauma patients: evaluation and management. *Hong Kong Med J*. 16(6):445-451.
8. Bauman MH, Strange C. (1992). Management of spontaneous pneumothorax. *Clin Chest Med*. 1992;13(4):645-658.
9. Macklin CC. (1939). Transport of air along sheaths of pulmonic blood vessels from alveoli to mediastinum: clinical implications. *Arch Intern Med*. 64:913-926.
10. American College of Chest Physicians. (2001). Management of spontaneous pneumothorax. *Chest*. 119(2):590-602.
11. Hernandez MC, Mora JA, Croce MA. (2019). Predictors of outcomes in trauma patients with rib fractures. *Am J Surg*. 218(5):983-989.
12. Singh G, Sharma A, Dhillon MS. (2018). Evaluation of traumatic pneumothorax: factors influencing outcomes. *J Emerg Med*. 54(4):475-480.

Ready to submit your research? Choose ClinicSearch and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At ClinicSearch, research is always in progress.

Learn more <https://clinicsearchonline.org/journals/clinical-research-and-studies->



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.