

Vaccination against SARS-CoV-2: Efficacy and Side effects

Rim M. Harfouch^{1*}, Lama Al Haushey², Yahya Elshimali³

¹Department of Microbiology, faculty of pharmacy, Alsham private university, Latakia, Syria, Department of biochemistry and microbiology, faculty of pharmacy, Tishreen university, Latakia, Syria

² Department of Pharmaceutics, faculty of pharmacy, Alsham private university, Latakia, Syria.

³Department of pathology, Faculty of medicine, Charles Drew University of medicine and science/ University of California Los Angeles (UCLA), USA.

***Corresponding Author:** Rim M. Harfouch. Department of Microbiology, faculty of pharmacy, Alsham private university, Latakia, Syria, Department of biochemistry and microbiology, faculty of pharmacy, Tishreen university, Latakia, Syria.

Received date: January 25, 2023; **Accepted date:** February 03, 2023; **Published date:** February 13, 2023.

Citation: Rim M. Harfouch, Lama Al Haushey, Yahya Elshimali. (2023). Vaccination against SARS-CoV-2: Efficacy and Side effects, *J. International Journal of Clinical Case Reports*, 2(1) DOI: 10.31579/2834-8389/006

Copyright: © 2023 Rim M. Harfouch. This is an open-access article distributed under the terms of The Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract:

The COVID-19 pandemic has spread rapidly and infected about 509 million people with 6.22 million deaths all around the world. Given the lack of specific therapy for and the rapid spread of this virus, vaccination would be a significant way in the fight against the SARS-CoV-2 pandemic. On December 11, 2020 and December 18, 2020, respectively, the US Food and Drug Administration (FDA), granted emergency authorization to the Pfizer/BioN-Tech and Moderna COVID-19 vaccines. In this review, we summarize the development of COVID-19 vaccine technology and define the approved vaccines by FDA. We also brief the most common side effects and differences observed between available vaccines.

Key words: COVID-19; vaccination; efficacy; side effects

Introduction:

The novel coronavirus SARS-CoV-2 or COVID-19 was first found in Wuhan, China and is the cause of severe acute respiratory distress syndrome. Afterwards, this virus spread rapidly and became a global pandemic [1]. Although the fatality rate is low (reported to be 2.5% as of 12 February 2020), the accelerating transmission makes it a threat to mankind, and finding a curative treatment is a top priority. While no such treatment has been confirmed, many drugs and combinations are being suggested and some have even shown positive clinical results. On January 23, 2020, the first clinical trial for COVID-19 was registered, the number of trials then ascended to reach 125 registered trials by February 18, 2020 [2].

Common symptoms of Covid-19 include acute respiratory illness (cold-like disease), hyperthermia (fever >38° C), coughing, sore throat, and shortness of breath. Also, numerous sufferers may also experience digestive symptoms such as anorexia, diarrhea, and vomiting [3].

During the COVID-19 pandemic, people are facing major health care challenges, lockdowns, and stress, as there is no specific treatment and vaccination for this pandemic. Given the lack of specific therapy for and the rapid spread of this virus, vaccination would be a significant way in the fight against the SARS-CoV-2 pandemic [4,5].

Development of COVID-19 vaccines:

Pfizer-BioNTech vaccine (BNT162b2) is based on the mRNA technology to express the SARS-CoV-2 spike (S) gene and has shown a high efficacy rate against SARS-CoV-2 infection. Specifically, phase III trials showed that BNT162b2 has about 95% efficacy against laboratory-confirmed SARS-CoV-2 symptomatic infection, at least seven days after the second dose in the individual of 16 years and older without current or previous history of COVID-19. mRNA vaccines are a new type of vaccine that has been recently utilized. BNT162b2 mRNA vaccine has been developed to stimulate immune response against SARS-CoV-2 using mRNA coding SARS-CoV-2 spike protein. This vaccine was approved by the U.S. food and drug administration (FDA) on the 11th of December 2020 for EUA in individuals older than 16 years of age. On the other hand, two doses of Oxford-AstraZeneca adenovirus-vectored vaccine (ChAdOx1 nCoV-19) showed an overall 63% efficacy against symptomatic SARS-CoV-2 infection. This vaccine was authorized to be used in the age group of 18 years and older. Unlike BNT162b2, ChAdOx1 nCoV-19 uses replication-deficient chimpanzees adenovirus as a viral vector to express the SARS-CoV-2 spike protein [6].

During 2020, 58 vaccines against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) be developed and in clinical trials, with some vaccines reportedly having more than 90% efficacy against COVID-19 in clinical trials. This remarkable achievement is much-needed good news as COVID-19 cases are currently at their highest daily levels globally. New vaccine efficacy results are reported now in *The Lancet*: investigators of four randomized, controlled trials conducted in

the UK, South Africa, and Brazil report pooled results of an interim analysis of safety and efficacy against COVID-19 of the Oxford–AstraZeneca chimpanzee adenovirus vectored vaccine ChAdOx1 nCoV-19 (AZD1222) in adults aged 18 years and older [7, 8].

On December 11, 2020 and December 18, 2020, respectively, the US Food and Drug Administration (FDA), granted emergency authorization to the Pfizer/BioN-Tech and Moderna COVID-19 vaccines. These two COVID-19 vaccines were developed quickly to benefit humanity and arrest the rise in the number of SARS-CoV-2 cases. From the time when the SARS-CoV-2 genome was released in early 2020 until these two

vaccines received EUA status, less than one year passed. The fastest any vaccine had previously been developed, from viral sampling to approval, was four years, for mumps in the 1960s. There have been some concerns about potential adverse effects of these vaccines. The present study aims to highlight evidence about the pharmacological characteristics, indications, contraindications and adverse effects of Pfizer/BioNTech and Moderna vaccines [9].

The following table (1) summarize the FDA approved COVID-19 vaccines with details on shots number, efficacy and side effects.

Vaccine developer	Pfizer	Moderna	AstraZeneca	Johnson & Johnson	Sputnik	Novavax
Mechanism	Messenger RNA	Messenger RNA	Inactivated cold virus	Modified cold virus	Modified Adenovirus	Stabilized form of the coronavirus spike protein
When approved/expected approval	Given full FDA approval Aug. 23, 2021	Dec. 18, 2021	May, 2021	Feb. 27, 2021	August 20, 2021	Not yet available.
Efficacy	95%	94.1%	70%	66.1% globally; 72% in the U.S.	90%	89.7%
Who is it recommended for?	People 5 years and older	People 18 years and older	Not yet available	People 18 years and older	People 18 years and older	Not yet available
How many shots do you need?	Two doses, 3 weeks apart	Two doses, 4 weeks apart	Two doses, a month apart	One dose	Two doses, 21 days apart	Two doses, 21 days apart
When might you become eligible for a booster shot?	At least 5 months after your primary COVID-19 vaccination series (for people 16 years and older).	At least 6 months after your primary COVID-19 vaccination (for people 18 years and older)	To be determined	At least 2 months after your primary J&J COVID-19 vaccination (for people 18 years and older)	At least 2 months after your primary shoot	
What are the side effects?	Fatigue, headache, chills, muscle pain, especially after the second dose	Fever, muscle aches, headaches lasting a few days. Effects worse after second dose.	Pain where you get the shot, fever, muscle aches, headache	Pain where you get the shot, headache, fatigue, muscle pain		Pain and tenderness, fatigue, headache, muscle pain
Any warnings?	The FDA issued a warning about heart inflammation. Since April 2021, there have been more than a thousand reports of myocarditis and pericarditis.	The FDA issued a warning about heart inflammation. Since April 2021, there have been more than a thousand reports of myocarditis and pericarditis.		The FDA issued a warning about an increased risk for developing Guillain-Barre syndrome.		
Any significant side effects?	Extremely rare cases of anaphylaxis in people who received the vaccine. Extremely rare cases of Bell's palsy, a type of temporary facial paralysis.	Extremely rare cases of anaphylaxis in people who received the vaccine. Extremely rare cases of Bell's palsy, a type of temporary facial paralysis.	Four total serious side effects, including two cases of transverse myelitis	There is a possible, rare relationship between this vaccine and blood clots with low platelets.	injection site pain, muscle pain, fatigue, fever, and headache	Not yet available
What about people with lowered immunity?	OK for people with HIV or immunosuppressing drugs.	OK for people with HIV or immunosuppressing drugs.	Not yet available	Not yet available		Not yet available

Table 1: the FDA approved COVID-19 vaccines

Side effects of covid-19 vaccines:

Clinical trials have shown that both Pfizer-BioNTech and Oxford-AstraZeneca vaccines were associated with various mild to moderate side effects, such as pain, redness or swelling at the site of injection, tiredness, headaches, chills, muscle, and joint aches, and fever.

According to a Saudi study, the participants who received the Oxford-AstraZeneca vaccine reported a significantly higher frequency of fatigue and headache than those who received the Pfizer-BioNTech vaccine [6].

Significant differences were observed between the side effect profiles of mRNA versus viral vector vaccines (predominantly Pfizer versus AstraZeneca). Overall, the recipients of mRNA vaccines reported a higher incidence of any self-reported side effects, which were, however, of significantly milder severity compared with those who received viral vector vaccines. While mRNA vaccines were associated with an increased incidence of local reactions, they were associated with a considerably lower incidence of systemic side effects including anaphylaxis, fever, swelling in the face or mouth or generalized swelling, flu-like illness, breathlessness and fatigue. Most importantly, mRNA vaccines were associated with a significantly lower incidence of severe side effects (requiring hospital care) [10].

On the other hand, another study conducted in Japan showed that individuals vaccinated with the mRNA-1273 vaccine were more likely to experience systemic reactions than those vaccinated with the BNT162b2 vaccine. Delayed injection site reaction was reported most frequently in middle-aged females after receiving the first dose of the mRNA-1273 vaccine [11].

The most common side effects of Sputnik V (adenovirus vector) are: injection site pain, fever, headache, fatigue, and muscle and joint pain. Moreover, unusual thrombotic cases have rarely been reported during the use of vector vaccines. The most important reasons involved in causing this rare complication are platelets and PF4 (platelet factor 4). The mechanisms that may be behind thrombotic thrombocytopenia after COVID-19 vaccination are: a) antibodies against PF4, b) the cross-reactivity of anti-spike antibodies and PF4, c) cross-reactivity of antibodies against adenovirus with PF4, d) interaction between spike protein and platelets, e) the direct interaction between adenoviral vector and platelets [12].

Conclusion:

The efficacy of FDA-approved vaccines against COVID-19 ranged from 70-95% with the highest efficacy of Pfizer-BioNTech vaccine. mRNA based vaccines (Pfizer and Moderna) may have lower systematic side effects compared with vector based vaccines (Sputnik V and AstraZeneca), but mRNA vaccines have higher incidence of local reactions and may be associated with myocarditis and pericarditis

according to thousand reports. Further studies on vaccine safety are recommended to strengthen public confidence in COVID-19 vaccines.

References:

1. Harfouch RM (2022). Antiviral Effects of Propolis against SARS-COV 2. *Int J Clin Med Imaging* 9:809
2. Harfouch, R.M., Alshaikh, S., Alshimaly, M., Assaad, A., Ahmad, J., Zoughaibi, H., Hammadi, M. and Elshimali, Y., (2021). Therapeutic approaches for covid 19: Challenges and successes. *Annals of Clinical and Analytical Medicine*, pp.228-233.
3. Jebur A , Harfouch T and Harfouch R. (2021) COVID-19 Pandemic in the Arab World: A Mini-Review Article. *CPQ Medicine*. 11(4), 01-10.
4. Harfouch RM. (2021) Cytokine Storm Syndrome in COVID-19 Patients: Characteristics and Diagnosis. *Ann Clin Med Case Rep*. 7(18):1-3.
5. Alhouri, Ali, et al. "Possible side effects of using detergents during the Covid19 pandemic in Syria." *Ann. Clin. Cases* 1.4 (2020): 1023.
6. Alhazmi A, Alamer E, Daws D, Hakami M, Darraj M, Abdelwahab S, Maghfuri A, (2021) Algaissi A. Evaluation of Side Effects Associated with COVID-19 Vaccines in Saudi Arabia. *Vaccines (Basel)*. 18;9(6):674
7. Knoll MD, Wonodi C. (2021) Oxford-AstraZeneca COVID-19 vaccine efficacy. *Lancet*. 397(10269):72-74.
8. Al-Salam, J.A., Ibrahim, H., and Shoujaa, A. (2022) "Pregnancy and COVID – 19: A Review of Tests, Medications and Vaccines." *Annals of pharma research* 10.2 Pp 662-665
9. Meo SA, Bukhari IA, Akram J, Meo AS, Klonoff DC. (2021) COVID-19 vaccines: comparison of biological, pharmacological characteristics and adverse effects of Pfizer/BioNTech and Moderna Vaccines. *Eur Rev Med Pharmacol Sci*. Feb;25(3):1663-1669.
10. Mathioudakis AG, Ghrew M, Ustianowski A, Ahmad S, Borrow R, Papavasileiou LP, Petrakis D, Bakerly ND. (2021) Self-Reported Real-World Safety and Reactogenicity of COVID-19 Vaccines: A Vaccine Recipient Survey. *Life (Basel)*. 17;11(3):249.
11. Kitagawa H, Kaiki Y, Sugiyama A, et al. (2022) Adverse reactions to the BNT162b2 and mRNA-1273 mRNA COVID-19 vaccines in Japan. *J Infect Chemother*. 28(4):576-581.
12. Zare H, Rezapour H, Mahmoodzadeh S, Fereidouni M. (2021) Prevalence of COVID-19 vaccines (Sputnik V, AZD-1222, and Covaxin) side effects among healthcare workers in Birjand city, Iran. *Int Immunopharmacol*. 101:108351.

Ready to submit your research? Choose ClinicSearch and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At ClinicSearch, research is always in progress.

Learn more <http://clinicsearchonline.org/journals/international-journal-of-clinical-case-reports->



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.