

Hypoglycaemia and Cardiac Failure: An Uncanny Association

Dweepala P.D.K. Priyadarshini, Mahismita Patro, Shweta Anand, R.S. Pal, Sameer Vaidya and Dipti Gothi *

Department of Pulmonary Medicine, ESI-PGIMSR, Basaidarapur, New Delhi, India.

*Corresponding Author: Dipti Gothi, Department of Pulmonary Medicine, ESI-PGIMSR, Basaidarapur, New Delhi, India.

Received date: November 26, 2024; Accepted date: December 09, 2024; Published date: December 27, 2024

Citation: Dweepala P.D.K. Priyadarshini, Mahismita Patro, Shweta Anand, R.S. Pal, Sameer Vaidya, (2024), Title, *International Journal of Clinical Case Reports*, 3(6); DOI:10.31579/2834-8389/034

Copyright: © 2024, Dipti Gothi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Department of Pulmonary MHypoglycaemia in the intensive care unit is frequently encountered with varied causes. Here we report an extremely rare case of persistent hypoglycaemia due to congestive cardiac failure (CCF) and its successful improvement with management of CCF. An 81 years old man with chronic kidney disease and coronary artery disease presented with recurrent syncope, dyspnoea and swelling of body to the emergency room. The detailed laboratory work-up revealed diagnosis of CCF. He was managed with intravenous dextrose infusion but the hypoglycaemia episodes were persistent and recurrent. The hypoglycaemia episodes were corrected only after improvement of CCF, thus confirming the cause of hypoglycaemia to be CCF. CCF has been reported to cause hypoglycaemia mostly in children but extremely rare in adults. CCF though rare should be considered as one of the causes of persistent hypoglycaemia in adults. Literature search revealed few case series implicating CCF as cause of hypoglycaemia.edicine, ESI-PGIMSR, Basaidarapur, New Delhi, India.

Keywords: hypoglycaemia; cardiac failure; uncanny association; congestive; diabetes mellitus; drug-induced; critical illnesses; organ failure

Introduction

Hypoglycemia is a common occurrence in the intensive care units (ICU). Persistent hypoglycaemia is one of the common end-stage manifestations in many diseases. The causes of hypoglycaemia include diabetes mellitus, drug-induced, critical illnesses like organ failure, certain hormonal deficiencies like cortisol or glucagon deficiency, non-islet cell tumour, malicious hypoglycaemia etc. The rare causes include congenital hyperinsulinism, insulin receptor mutation, inborn errors of metabolism, ectopic insulin secretion, autoimmune causes, cardiac failure etc [1-3]. Here we are reporting one such case of hypoglycemia due to congestive cardiac failure (CCF) which is of extremely rare occurrence.

Case Report

An 81-year-old man, never smoker presented to the emergency room (ER) with altered sensorium, decreased food intake and irrelevant talking for two days. He had no history of fever, cough or shortness of breath. He had coronary artery disease (CAD) with history of percutaneous transluminal coronary angioplasty with stenting to left anterior descending artery and right coronary artery two months ago. His echocardiography done at that time showed a left ventricular ejection fraction of 25-30% with moderate mitral regurgitation and trivial tricuspid regurgitation. He was also diagnosed with chronic kidney disease (CKD) 3 months back. His prior blood urea was 137 mg/dl and serum creatinine were 3.5 mg/dl and he was on conservative management for CKD. There were no other comorbidities. At presentation in the ER, he was conscious but confused with Glasgow coma scale score of 12. His respiratory rate was 26/minute, pulse rate was 94/minute, oxygen saturation was 92% on room air and blood pressure was 140/90 mmHg. On auscultation, there were normal vesicular breath sounds and bilateral fine basal crepitations. The heart sounds were normal.

His random blood sugar (RBS) in the ER was 35 mg/dl. The arterial blood gas analysis showed pH of 7.49, pCO₂ of 26.8,

pO₂ of 52, and HCO₃ of 20.9. His electrocardiogram had a normal sinus rhythm and troponin-T was negative. His chest radiograph showed cardiomegaly with bilateral middle and lower zone opacities in perihilar predominance indicating pulmonary oedema. He was given intravenous dextrose for correction of hypoglycaemia and then shifted to ICU with the provisional diagnosis of CCF with pulmonary edema vs bilateral pneumonia with sepsis and hypoglycaemia.

In the ICU he had similar episodes of hypoglycaemia which were recurrent and was managed by continuous infusions of 25

Discussion

Hypoglycaemia is defined as a plasma glucose concentration below 4 mmol/L (72 mg/dl) [4]. Occurrence of episodes of hypoglycaemia in critically ill patients is known [5]. It occurs due to septic shock related adrenal failure or fulminant hepatic failure. The cause of hypoglycaemia in critically ill patients is combination of insulin excess and inadequate nutritional supplementation along with decreased glucose production endogenously and increased glucose utilisation [5-7].

CCF is one of the rare and under-recognised causes of hypoglycaemia [8]. The mechanism of hypoglycaemia in CCF is poorly understood. Glucose and fatty acids are the main energy source for the myocardial cells [9]. Myocardial tissue preferentially uses glucose as the main source of energy in case of dilated or hypertrophic cardiomyopathy and during hypoxia or ischemia [10]. Hypoglycaemia induced plasma glucose fluctuations enhances reactive oxygen species production and exaggerated myocardial injury further leading to worsening of myocardial function [11,12]. The mechanism of hypoglycaemia in CCF is through the hepatic involvement. Liver injury in heart

failure causes glycogenesis and gluconeogenesis leading to hypoglycaemia. Low calorie intake, increased glucose utilization by ischaemic tissues including the heart, insufficient peripheral circulation and malabsorption of nutrients due to gastrointestinal oedema are other possible mechanisms of hypoglycaemia in heart failure patients [13-15].

The clinical presentation of hypoglycaemia ranges from dizziness, sweating, palpitations, tremor, weakness, confusion, altered mental status to life-threatening complications like seizures and coma [4]. Immediate diagnosis and prompt glucose infusions is the key to successful treatment of hypoglycaemia and prevention of life-threatening complications. Prompt diagnostic approach towards the cause of the hypoglycaemia is necessary to prevent further such episodes.

Spontaneous hypoglycaemia associated with CCF is rarely reported in adults, though more cases have been reported in neonates and infants. **In a case report by H Khoury et al.**, an inverse relationship was found between serum insulin level and serum glucose level [16]. Hypoglycaemia resolved after alleviation of the CCF. In a study conducted by Benzing et al., acute CCF and hypoglycaemia were observed simultaneously in 27 patients with underlying heart disease and also found that constant intravenous infusion of glucose is the method of choice for the treatment of these patients rather than epinephrine or glucagon along with treatment of heart failure [15]. Hedayat et al., reported eleven cases of spontaneous hypoglycaemia in CCF in adults [13].

All the reported cases have found an association between cardiac failure and persistent hypoglycemia [13-17]. To the best of our knowledge this is the second case from India showing this interesting and important association of hypoglycaemia and CCF [17]. One recent clinical study in 2020 conducted by Y Teshima et al., found out that patients with chronic heart failure are at risk of developing hypoglycaemia even if they do not undergo any antidiabetic therapy [18]. It is possible that digging into the cause of hypoglycaemia is not given priority rather than its treatment. And even if found they may not be reported since there is paucity of evidence and knowledge regarding this association. Thus, we want to highlight that it is pertinent to find out the cause of hypoglycaemia in each case.

CKD is also an unusual cause of persistent spontaneous hypoglycaemia like CCF [19]. In CKD there is reduced glomerular filtration rate which leads to reduced clearance of insulin hence causes hypoglycaemia [20]. Our patient had both CKD and CCF. But in our case, we concluded that the cause of persistent hypoglycaemia as CCF as the hypoglycaemia resolved with the successful treatment of CCF and our patient was only on conservative management for his renal disease without any dialysis or acute management.

Conclusion

CCF should be considered as one of the causes of persistent spontaneous hypoglycaemia and thoroughly investigated. Timely suspicion and management can improve the outcomes by prevention of further episodes of hypoglycaemia and worsening of cardiac function by repeated episodes of hypoglycaemia.

References

- Effect of intensive diabetes treatment on the development and progression of long-term complications in adolescents with insulin-dependent diabetes mellitus: Diabetes Control and Complications Trial. Diabetes Control and Complications Trial Research Group. *J Pediatr.* 1994 Aug;125(2):177–188.
- Cryer PE. Hypoglycaemia: the limiting factor in the glycaemic management of Type I and Type II diabetes. *Diabetologia.* 2002 Jul;45(7):937–948.
- Douillard C, Jannin A, Vantghem MC. Rare causes of hypoglycemia in adults. *Ann Endocrinol.* 2020 Jun 1;81(2):110–117.
- Yale JF, Paty B, Senior PA. Hypoglycemia. *Can J Diabetes.* 2018 Apr;42:S104–108.
- Pratiwi C, Mokoagow MI, Made Kshanti IA, Soewondo P. 2020 The risk factors of inpatient hypoglycemia: A systematic review. *Heliyon.* May 1;6(5): e03913.
- Cryer PE. Hypoglycaemia: 2006 the limiting factor in the glycaemic management of the critically ill? *Diabetologia.* Aug;49(8):1722–1725.
- Lacherade JC, Jacqueminet S, Preiser JC. An Overview of Hypoglycemia in the Critically Ill. *J Diabetes Sci Technol.* 2009 Nov 1;3: 1242–1249.
- [Bansal N, Weinstock RS. Non-Diabetic Hypoglycemia. In: 2000 Feingold KR, Anawalt B, Boyce A, Chrousos G, de Herder WW, Dhatariya K, et al., editors. Endotext \[Internet\]. South Dartmouth \(MA\): MDText.com, Inc.](#)
- Lopaschuk GD, Ussher JR. Evolving Concepts of Myocardial Energy Metabolism. *Circ Res.* 2016 Nov 11;119(11):1173–1176.
- Ardehali H, Sabbah HN, Burke MA, Sarma S, Liu PP, Cleland JGF, et al. Targeting myocardial substrate metabolism in heart failure: potential for new therapies. *Eur J Heart Fail.* 2012 Feb;14(2):120–129.
- Saito S, Thuc LC, Teshima Y, Nakada C, Nishio S, Kondo H, et al. Glucose Fluctuations Aggravate Cardiac Susceptibility to Ischemia/Reperfusion Injury by Modulating MicroRNAs Expression. *Circ J Off J Jpn Circ Soc.* 2016;80(1):186–195.
- Snell-Bergeon JK, Wadwa RP. Hypoglycemia, Diabetes, and Cardiovascular Disease. *Diabetes Technol Ther.* 2012 Jun;14(Suppl 1):S-51-S-58.
- Hedayati HA, Beheshti M. Profound spontaneous hypoglycaemia in congestive heart failure. *Curr Med Res Opin.* 1977;4(7):501–504.
- Block MB, Gambetta M, Resnekov L, Rubenstein AH. Spontaneous hypoglycaemia in congestive heart-failure. *Lancet Lond Engl.* 1972 Oct 7;2(7780):736–738.
- Benzing G, Schubert W, Sug G, Kaplan S. Simultaneous hypoglycemia and acute congestive heart failure. *Circulation.* 1969 Aug;40(2):209–216.
- Khouri H, Daugherty T, Ehsanipoor K. Spontaneous hypoglycemia associated with congestive heart failure attributable to hyperinsulinism. *Endocr Pract Off J Am Coll Endocrinol Am Assoc Clin Endocrinol.* 1998 Apr;4(2):94–95.
- Yawar Y, Ahmad SP, Irfan M, Saika A. A Rare Case Of Hypoglycemic Unawareness In A Patient With Chronic Congestive Heart Failure. 2019 Jan 1;4(1):20–22.
- [Potential Risk of Hypoglycemia in Patients with Heart Failure \[Internet\]. \[cited 2022 Jul 21\].](#)
- Moen MF, Zhan M, Hsu VD, Walker LD, Einhorn LM, Seliger SL, et al. Frequency of hypoglycemia and its significance in chronic kidney disease. *Clin J Am Soc Nephrol CJASN.* 2009 Jun;4(6):1121–1127.
- Ahmad I, Zelnick LR, Batacchi Z, Robinson N, Dighe A, Manski-Nankervis JAE, et al. Hypoglycemia in People with Type 2 Diabetes and CKD. *Clin J Am Soc Nephrol.* 2019 Jun 7;14(6):844–853.