

The Efficacy of Dzarirah oil (*Acorus calamus* L.) Against Bacteria *Propionibacterium Acnes* that Induce Maskne Disease

Amalia Tri Utami^{1*}, Ghassan Ahmed¹ Abdullah Qarah²

¹Faculty of Medicine and Health, *State University of Malang*, Indonesia

²Faculty of Medicine and Medical Science, University of Aden, Yemen

*Correspondence Author: Amalia Tri Utami, Faculty of Medicine and Health, State University of Malang, Indonesia

Received Date: March 03 2023 | Accepted Date: March 17 2023 | Published Date: March 30 2023

Citation: Amalia Tri Utami, Ghassan Ahmed Abdullah Qarah (2023). The Efficacy of Dzarirah oil (*Acorus calamus* L.) Against Bacteria *Propionibacterium Acnes* that Induce Maskne Disease. *International Journal of Clinical Case Studies*.2(2); DOI:10.31579/ijccs-2022/006

Copyright: © 2023 Amalia Tri Utami, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Maskne is a pimple that occurs as a result of using a mask with hot ingredients and using it for too long. Maskne is a new term that comes from the word mask plus acne. According to dermatologists, the continuous use of masks causes a build-up of bacteria and dirt that clogs the skin pores and triggers acne and other skin problems. The reason is, the skin area around the cheeks and chin does not get air circulation because it is covered by a mask. The Centers for Disease Control and Prevention and the World Health Organization require the use of masks for all people around the world to prevent exposure to the Corona virus which causes COVID-19 infection. However, there are effects from using this mask in the form of pimples and other facial skin irritations. The Maskne triggered by infection of *Propionibacterium acnes* which plays a role in irritation of the follicular epithelium and facilitates the occurrence of Maskne. Medical treatment is mostly done is by giving topical and oral antibiotics which are relatively more expensive and can cause side effects. As an alternative medicine in the community, dzarirah (*Acorus calamus* L.) can be used which has been scientifically proven to have antibacterial activity. Dzarirah is a type of plant that is mentioned in the hadith to treat the acne of the wife of Rasulullah SAW. The prophet's wives told that one day the Prophet met them. Rasulullah then saw among them a kind of pimple grew. Rasulullah then asked, "Do you have the perfume of dzarirah ?." They said they had. Rasulullah said, "Put your pimples on while reading a prayer, 'O Allah, who reduce the large and the small ones, reduce my acne'." The α and β asarone compounds are antimicrobial compounds that can inhibit the growth of *P. acnes* bacteria. The results showed that dzarirah oil could inhibit the growth of *Propionibacterium acnes* bacteria at a concentration of 2.5% while killing at a concentration of 25%. Dzarirah oil is effective in inhibiting the growth of *Propionibacterium acnes* bacteria at a concentration of 2.5% and is a MIC (Minimum Inhibitory Concentration). Dzarirah oil is proven to be effective in killing *Propionibacterium acnes* bacteria at a concentration of 25% and is an MBC (Minimum Bactericidal Concentration).

Keywords: dzarirah oil; *acorus calamus*; *propionibacterium acnes*; maskne

Introduction

Acorus calamus L. or also known as the sweet flag is a species in the Acoraceae family. *Acorus calamus* is a plant that has many benefits including medicine (Silalahi & Nisyawati 2018), foodstuff (Dušek et al. 2007), and is also useful in land conservation. Pai and McCarthy (2010) stated that *Acorus calamus* is one of the socio-economically important plant species used in wetland restoration projects that can act as an economic incentive for conservation and restoration.

At first, *Acorus calamus* L. was included in the Araceae family (taro), but eventually it was expelled from that family and included in the Arecaceae family because it had very different characteristics, especially in its leaf

structure and rhizome. Chang (2010) stated that the genus *Acorus* is not related to the Araceae family.

Acorus calamus L. has long been used by local Indonesian people as a fever medicine by the Simalungun Batak ethnic group (Silalahi et al. 2015a), and the Karo Batak (Silalahi et al. 2015b; Silalahi and Nisyawati 2018; Silalahi 2016), malnutrition (Silalahi et al. 2015b) and medicine for postpartum mothers by the Minangkabau ethnicity (Khariah et al. 2017). The many benefits of *Acorus calamus* have resulted in this plant being cultivated by the Angkola Mandailing Batak ethnic group in their yards (Silalahi 2016). Rhizoma is a part that is used in a single form or a potion.

The efficacy of plants as medicinal substances is related to their secondary metabolites. From various research reports, although rhizomes are the

most frequently used, almost all organs (roots, rhizomes, and leaves) of *A. calamus* contain essential oil. Essential oil is a compound of the terpenoid group, especially monoterpenoids and sesquiterpenoids which are volatile at room temperature (Harbone 1987). This results in plants that have essential oils to produce distinctive aromas (Dušek et al. 2007). Asaron is the main essence found in *A. calamus* (Hanelt et al. 2001). However, the essential oil content in *A. calamus* is strongly influenced by internal and external factors. This causes the essential oil content to differ from one region to another. Internal factors that affect the asaron content of the *Acorus calamus* L., especially the karyotype of the chromosome number (Hanelt et al. 2001). Although there have been many studies on *A. calamus*, comprehensive discussion of its use and secondary metabolite content is still very limited, especially its use as an anti-cancer and anti-microbial.

Acne is caused by several factors, one of which is the *Propionibacterium acnes* bacteria. Gaspari and Stephen (2008) stated that *Propionibacterium acnes* includes normal flora of the skin, gram-positive, pleomorphic, and anaerobic bacteria. These bacteria play a role in the formation of acne, by producing lipases that break down free fatty acids from skin lipids, causing inflammation. The resulting inflammation causes *Propionibacterium acnes* to proliferate and aggravates inflammatory lesions by stimulating the production of proinflammatory cytokines. These bacteria are not pathogenic under normal conditions, but if there is a change in skin conditions, the bacteria will turn out to be invasive. The secretions of sweat glands and sebaceous glands which produce water, amino acids, urea, salts, and fatty acids are a source of nutrition for

bacteria. These bacteria play a role in the chemotactic inflammation process and the formation of lipolytic enzymes that convert the oily substance fraction (sebum) into a solid mass, which causes blockage of the sebaceous gland ducts (Djuanda, et al. 1999). *Propionibacterium acnes* is a dangerous bacteria that can cause inflammation of the skin, especially on facial skin, so it needs to be prevented.

Method And Results

This research is a laboratory experimental study, using a completely randomized design (CRD) with 5 treatments and 4 replications. The techniques used are well, dilution techniques and colony counting techniques. The parameters in this study were the diameter of the bacterial growth inhibition zone, the color change formed in each bacterial dilution tube, and the calculation of the number of *Propionibacterium acnes* bacterial colonies.

The dilution method is divided into 2, namely liquid and solid dilution. The dilution method is used to measure the MIC (minimum inhibitory level) while the solid dilution method is used to determine the MBC (minimum bactericidal level). The method used in the liquid dilution method is to make a series of dilutions of the antimicrobial agent in liquid media which is added with the test microbes. The solid dilution method is carried out by inoculating the test microbes on agar media containing antimicrobial agents. The advantage of this dilution method is that the only antimicrobial control agent tested can be used for testing several test microbes (Pratiwi, 2008). The results of testing Dzarirah oil against *P. Acne* are presented in Figure 1.




Figure 1: Dzarirah oil dilution tube against *P. Acne*

The diffusion method was used to determine the sensitivity of the tested microbes to antimicrobial agents. This method is done using disc paper. A disc paper was inserted into the agar medium that had been inoculated with the bacteria and filled with the test compound. The clear area on the surface of the agar media indicates an inhibition of the growth of microorganisms by antimicrobial agents. The advantage of this diffusion method is that it is easy to do because it does not have special tools and includes greater flexibility in choosing the drug to be examined (Katrin et al., 2015).

The dilution method was used to determine the MIC and MBC of antimicrobial drugs. The principle of this dilution method is to use a series of test tubes filled with liquid media and *P. Acne* test bacteria cells. The

MIC of the drug is shown from the lowest concentration of antibiotics in the tube with the results of the culture starting to appear clear (no microbial growth). The lowest concentration of drug in a solid culture is indicated by the absence of growth of microbial colonies in the MBC of the drug against the tested bacteria. This experiment uses antibiotics from Dzarirah oil. In Table 1, the best value for teaching and learning activities in dzarirah oil is 25%.

Then done counting the number of colonies of *P. Acne* at each concentration with four repetitions by using Colony Counter. Counting the number of colonies of *P. Acne* at each Mc Conkey Agar. Results planting MAC media of various concentrations of the extract can be seen in Table 1

Treatment	Repetition				Mean
	1	2	3	4	
1.25%	2.1×10^7	2.2×10^7	2.3×10^7	3.4×10^7	2.5×10^7
1.5%	1.4×10^6	2.7×10^6	3.4×10^6	2.9×10^6	2.6×10^6
1.75%	4.7×10^5	4.2×10^5	4.1×10^5	4.2×10^5	4.3×10^5
2%	2.7×10^4	2.2×10^4	3.4×10^4	3.7×10^4	3×10^4
2.25%	1.2×10^3	1.3×10^3	1.7×10^3	1.9×10^3	1.5×10^3
2.5%	9.2×10^2	9.8×10^2	9.7×10^2	9.6×10^2	9.6×10^2
10%	4.2×10^1	4.5×10^1	6.2×10^1	7.2×10^1	5.5×10^1
25%	0%	0%	0%	0%	0%
50%	0%	0%	0%	0%	0%

Table 1. The Results Mean Colony of *P. Acne* in Different Treatment

Result

The α and β asarone compounds are antimicrobial compounds that can inhibit the growth of *P. acne* bacteria. The results showed that dzarirah oil could inhibit the growth of *Propionibacterium acnes* bacteria at a concentration of 2.5% while killing at a concentration of 25%. Dzarirah oil is effective in inhibiting the growth of *Propionibacterium acnes* bacteria at a concentration of 2.5% and is a MIC (Minimum Inhibitory Concentration). Dzarirah oil is proven to be effective in killing *Propionibacterium acnes* bacteria at a concentration of 25% and is an MBC (Minimum Bactericidal Concentration).

References

1. Chang, P.S. (2010). *Flora of China*. 23:1-2.
2. Djuanda, A., Hamzah, M. dan Aisah, S. (1999). *Ilmu Penyakit Kulit dan Kelamin*. Jakarta: Balai Penerbit FKUI.
3. Dušek, K., Galambosi, B., Hethelyi, E.B., Korany, K., & Karlová, K. (2004). Morphological and chemical variations of sweet flag (*Acorus calamus* L.) in the Czech and Finnish gene bank collection. *Hort. Sci. (Prague)* 34 (1): 17-25.
4. Khairiah, A. (2017). *Etnomedisin dan Nilai Ekonomi Tumbuhan Obat Pada Etnis Minangkabau di Kecamatan IX Koto Sungai Lasi, Solok, Sumatra Barat*. Tesis. Departement Biologi, FMIPA, Universitas Indonesia, Depok. Indonesia.
5. Gaspari, A.A and Stephen, K.T. (Eds) (2008). *Clinical and Basic Immunodermatology*. London: British Library Cataloging in Publication.
6. Hanelt P., (2001). *Mansfeld's Encyclopaedia of Agricultural and Horticultural Crops*. 5th vol., 1st ed. Berlin, Springer: 2317-2318.
7. Harborne, J.B. (1987). *Metode fitokimia: Penuntun cara modern menganalisis tumbuhan*, Ed. II. Diterjemahkan oleh Kosasih Padmawinata dan Iwang Soedira. ITB Press, Bandung: x + 354 hlm.
8. Katrin, D., Idawati, N. and Sitorus, B., (2015). Uji Aktivitas Antibakteri Dari Ekstrak Daun Malek (*Litsea gracie* Vidal) Terhadap Bakteri *Staphylococcus aureus* dan *Escherichia coli*. *Jurnal Kimia Khatulistiwa*, 4(1), pp.7-12.
9. Pratiwi, S.T., 2008. *Mikrobiologi Farmasi*, Jakarta: Erlangga.
10. Silalahi, M., Nisyawati, Walujo, E.B., & Supriatna, J. (2015). Local knowledge of medicinal plants in subethnic Batak Simalungun of North Sumatra, Indonesia, *Biodiversitas* 16(1), 44-54.
11. Silalahi, M., Nisyawati, Walujo, E.B., Supriatna, J. & Mangunwardoyo, W. (2015). The local knowledge of medicinal plants trader and diversity of medicinal plants in the Kabanjahe traditional market, North Sumatra, Indonesia. *Journal of Ethnopharmacology* 175, 432-443.
12. Silalahi, M. (2016). Diversity of medicinal plants in home gardens in Tanjung Julu village, North Sumatra, Indonesia. *International Journal of Biological Research Volume* 4(1): 78-82.
13. Silalahi, M. & Nisyawati. (2018). The ethnobotanical study of edible and medicinal plants in the home garden of Batak Karo sub-ethnic in North Sumatra, Indonesia. *Biodiversity* 19(1): 229- 238.
14. Pai, A. & McCarthy, B.C. (2010). Suitability of the Medicinal Plant, *Acorus calamus* L., for Wetland Restoration. *Natural Areas Journal* 30(4): 380-386.

Ready to submit your research? Choose ClinicSearch and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At ClinicSearch, research is always in progress.

Learn more <http://clinicsearchonline.org/journals/international-journal-of-clinical-reports-and-studies>



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.