

# Kinetics of Changes in Biochemical Parameters and Activity of the Blood lipase Enzyme in Partridges (*Alectoris chukar*) with the Complex use of Vitamins and an Anti-Stress Drug

Topchiyeva Sh. A \*

Institute of Zoology of Ministry of Science and Education of the Republic of Azerbaijan

\*Correspondence Author: Topchiyeva Sh. A, Institute of Zoology of Ministry of Science and Education of the Republic of Azerbaijan.

Received Date: July 13 2023 | Accepted Date: August 04 2023 | Published Date: August 21 2023

**Citation:** Topchiyeva Sh. A, (2023), Kinetics of Changes in Biochemical Parameters and Activity of the Blood lipase Enzyme in Partridges (*Alectoris chukar*) with the Complex use of Vitamins and an Anti-Stress Drug, *International Journal of Clinical Case Studies*,2(4); DOI:10.31579/2835-9291/010

**Copyright:** © 2023, Topchiyeva Sh. A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract:

The aim of this work was to study the activity of the lipase enzyme in the blood serum of partridges. In experiments, experimental groups of partridges were fed with a complex of vitamins (A-20000 IU, D3 - 1250 IU, E - 50 mg) and an anti-stress drug (succinic acid) at a dose of 0.01, 0.03, 0.05 g per 1 kg of body weight per day. Determination of the activity of the lipase enzyme in the blood serum of partridges was carried out on a Specol 1500 spectrophotometer (Analitik Jena) according to the method of R Rej., M Hoder, U.H Bergmeyer at a wavelength of 540 nm. At the same time, the activity of lipase in the blood of control partridges was within the normal range (11.2-26.2 U/l). There were no statistically significant differences between the groups ( $P > 0.05$ ). As a result of experimental studies, the activity of the enzyme lipase in the blood serum of the experimental groups of partridges was revealed, which was 1.02, 1.56, 1.30, 1.29, 1.15, 1.13, 1.16, 1.24, 1.53, 1.29 times more, compared with the indicators of the enzymatic activity of the control groups of partridges. The level of lipase activity in partridges was experimentally revealed with the complex use of vitamins (A, D3, E) and the anti-stress preparation succinic acid, which reached  $11.5 \pm 0.7$  U/l in one-day-old partridges, and a maximum value of  $33.8 \pm 4.7$  U/l in 30- daily. Thus, the results of experimental studies can be used for scientific purposes to study the influence of various factors on the biochemical parameters of bird blood.

**Keywords:** enzyme activity; lipase; blood; partridges; *Alectoris chukar*; vitamins; anti-stress drug

## Introduction

Poultry farming is considered one of the actively developing branches of agriculture, the main condition for growth is the use of feed resources and the proper organization of full-fledged feeding of poultry. The most important is the provision of diets with protein and vitamin supplements, as well as enzymes, vitamins, amino acids, antibiotics and other biologically active substances that increase productivity and accelerate the growth of poultry.

Poultry farming provides the country's population with such highly nutritious and dietary products as eggs and meat, supplies light industry with down and feathers, and agriculture provides organic fertilizers. In this regard, agricultural birds deservedly enjoy increased attention from scientists of various specialties.

The literature available to us covers mainly the issues of cultivation technology, maintenance, frequency of feeding; composition and type of feed, egg productivity of the studied birds.

Also, to assess the effect on the metabolism of chickens, an experiment was conducted in which the features of the action of the feed additive Bacell were studied, depending on the dosage of its introduction into the feed of broiler

chickens of the "SK Rus-6" cross. For what formed four experimental groups of daily broilers with 56 heads each. Group 1 chickens that received no supplements served as controls.

The activities of pancreatic enzymes in pancreatic juice, blood serum, and pancreatic homogenate (in broilers) were determined using the kinetic methods. Activity of amylase in pancreatic juice and pancreatic homogenate was determined using Smith-Roy method modified by Ugolev [2] and expressed as mg of hydrolyzed starch per 1 ml of juice (or 1 g of homogenate) per min; activity of lipase on semi-automatic biochemical analyzer BS3000P (China) using reagent kit by DIAKON-VET Co. (Russia) and expressed as units per 1 ml of juice (or 1 g of homogenate). The activities of amylase and lipase in blood serum were determined on automatic biochemical analyzer Chem Well 2900 (T) (USA) using reagent kits Liquicolor (Pancreas-Amylase and Lipase, Human, Germany) and filters 405 nm for amylase and 580 nm for lipase; the results were expressed as units per liter [3].

Activity of proteases in pancreatic juice and pancreatic homogenate was determined by casein degradation with colorimetric control [4] and expressed as mg of hydrolyzed casein per 1 ml of juice (or 1 g of

homogenate) per min. The activity of trypsin in blood serum were determined on semi-automatic biochemical analyzer BS3000P (China) using BAPNA (N- $\alpha$ -Benzoyl-D, L-arginine 4-nitroanilide hydrochloride) as a substrate and expressed as units per liter.

The activities of enzymes in pancreatic juice and blood serum can be influenced by different factors. Ingestion of feed and its movement down the gastro-intestinal tract is a powerful stimulator of exocrine pancreatic secretion; during the first minutes after the feeding amounts of juice and enzymes within the juice rise substantially compared to the corresponding pre-prandial levels (in starved birds) due to the complex-reflex phase of regulation [5].

Animal fats and vegetable oils are being used in broiler diets to increase energy density and improve growth performance [7,8]. However, fat addition negatively affects fat digestibility [9,10] especially during early broiler age [11]. Immature physiological function of the pancreas in broilers results in less production of bile acids and pancreatic lipase during early ages [12], which may lead to poor fat digestibility.

Supplementation of bile acids in broiler diet significantly improve the digestibility of fat [13]. Similarly, other studies [14,15,16] reported that the supplementation of bile acids improve daily weight gain, feed conversion ratio (FCR) and carcass yield in broilers. Like bile acids, exogenous lipase also improves physiological limitation of poultry digestive system [17]. On the other hand, Hu et al. [18] reported that providing reduced energy diet had decreased ( $p < 0.05$ ) body weight (BW) gain compared to basal energy diet during a period of 14 days, however, it was compensated with the supplementation of 0.015% and 0.03% lipase. According to Wang et al. [19], the supplementation of lipase in broiler diets improved FCR, growth performance and fat digestibility. On the contrary, other researchers reported that lipase supplementation had no effect on nutrient utilization and bird's performance in broiler fed wheat-based diets [20].

The literature data obtained make it possible to objectively assess the relationship of metabolic processes in the organism of the studied birds and their influence on the level of activity of blood enzymes. However, the given data of scientific sources do not give a complete picture and many aspects need a deeper study. Based on the foregoing, the aim of this work was to study the activity of the lipase enzyme in the blood of partridges.

## Material and research methods:


Experimental studies on the activity of the lipase enzyme in the blood serum of partridges were carried out on 35 birds, which were divided into control and experimental groups of 5 birds for 120 days. The control group of partridges were kept under standard conditions without the addition of vitamins and an anti-stress drug to their diet. During feeding, experimental groups of partridges were fed with a complex of vitamins (A-20000 IU, D3 - 1250 IU, E - 50 mg) and an anti-stress drug (succinic acid) at a dose of 0.01, 0.03, 0.05 g per 1 kg of body weight per day.

Determination of the activity of the lipase enzyme in the blood of partridges was carried out on a Specol 1500 spectrophotometer (Analytik Jena) according to the method of R Rej., M Hoder, U.H Bergmeyer at a wavelength of 540 nm [21].

The experimental groups of partridges were divided into three groups: the 1st experimental group was given a vitamin complex, the 2nd experimental group was given an anti-stress drug, the 3rd experimental group was added a complex of vitamins and an anti-stress drug. To determine the activity of the lipase enzyme in the blood serum of partridges, blood was obtained from the axillary vein before feeding and one hour after feeding. A heparin solution was added to the tubes, the blood was centrifuged at 5000 rpm for 5 min, followed by determination of the enzyme activity. Blood serum was examined for lipase activity. Statistical processing of the research results was performed using the Excel computer program, determining the mean value (M) and standard errors of the mean (m). Significance of differences was assessed by Student's t-test. Differences were considered statistically significant at  $p < 0.05$ .

## Research results:

As a result of our experimental studies, it was found that the activity of lipase in the blood serum of both control and experimental groups of partridges was undulating with a maximum in the control group by the age of 5, 10, 18 days and a minimum at the age of 1, 30 days, and in experimental groups of partridges with a maximum of 3, 10, 30, and with a minimum activity of 1, 5, 15 days of age, which, in all likelihood, is associated with a change in diet. The enzymatic activity of lipase in the blood of experimental groups of partridges, which simultaneously included in the diet both a complex of vitamins and an anti-stress drug, was more overestimated than the experimental groups, in which only vitamins or only an anti-stress drug were separately included in the diet, reaching a maximum value by 30 days (figure.1).



**Figure 1: Diagram of the effect of vitamins and an anti-stress drug on the activity of pheasant blood serum lipase.**

The activity of the lipase enzyme in the blood serum of the experimental groups of partridges, when a complex of vitamins and an anti-stress drug was included in the diet, increased by 1,3,5,7,10,15,18,20,25 30 days at 1.02,

1.56, 1.30, 1.29, 1.15, 1.13, 1.16, 1.24, 1.53, 1.29 times compared to lipase activity in partridge control groups (tabl.1).

During the period from 1 to 30 days of partridge, which were fed with a complex of vitamins (A-20000 IU, D3 - 1250 IU, E - 50 mg) and an anti-stress drug (succinic acid) at a dose of 0.01, 0.03, 0.05 g per 1 kg of body weight per day had a higher enzymatic activity compared to the control groups of birds. Partridges of the experimental groups, in which only

vitamins or only an anti-stress drug were added to the diet, differed in lower indicators of lipase activity than the experimental groups of birds, in which a complex of vitamins and an anti-stress drug were included in the diet. At the same time, lipase activity was higher ( $p < 0.05$ ) than with separate addition of a vitamin complex or an anti-stress drug to the diet.

Age in days	Control	1 experienced	2 experienced	3 experienced
1	11.2±0.2	11.4±0.3	11.3±0.9	11.5±0.7
3	12.6±2.1	19.5±0.9	18.3±1.6	19.7±0.3
5	16.1±1.3	20.6±1.2	19.5±0.8	20.9±0.2
7	14.3±1.1	18.1±2.1	17.5±1.7	18.4±1.7
10	20.0±0.9	22.8±0.9	21.3±1.6	23.0±2.8
15	18.1±1.6	19.2±1.3	18.7±2.8	20.5±3.1
18	24.5±3.1	27.1±4.1	24.7±3.1	28.2±3.5
20	21.3±2.6	25.1±3.4	23.6±3.1	26.4±4.0
25	16.5±2.1	24.3±5.1	23.1±3.4	25.3±5.1
30	26.2±0.7	32.1±4.7	30.5±5.8	33.8±4.7

**Table 1: Influence of vitamins and anti-stress drug on the activity of partridge blood serum lipase (U/l) ( $n=5$ ,  $M \pm m$ )**

By the activity of lipase, we can judge the state of the pancreas. As can be seen from the data in the table, the lipase activity of the blood serum of control partridges was within the normal range (11.2-26.2 U/l). There were no statistically significant differences between the groups ( $P > 0.05$ ).

Thus, the experimental data obtained by us are of a fundamental nature and can be used for scientific purposes to study the influence of various factors (physical, chemical, biological) on the activity of enzymes.

It should be noted that a properly composed diet for birds is no less important than their genetic origin. Therefore, it is necessary to take into account not only the nutritional value of the feed ration, but also the presence of biologically active substances in them.

With the rational and purposeful use of biologically active additives in the diets of birds, it has a positive effect not only on metabolic processes in the body, but also with the positive effect of the drugs used, productivity increases and the profitability of poultry products increases.

## Conclusions:

1. The activity of the enzyme lipase in the blood serum of the experimental groups of partridges was established, which was 1.02, 1.56, 1.30, 1.29, 1.15, 1.13, 1.16, 1.24, 1.53, 1.29 times higher, compared with the indicators of the enzymatic activity of the control groups of partridges.
2. The level of lipase activity in partridges was revealed with the complex use of vitamins (A, D3, E) and the anti-stress preparation succinic acid, which reached a value of  $11.5 \pm 0.7$  U/l in partridges, and a maximum value of  $33.8 \pm 4.7$  U/l in 30 -day.

## Reference:

1. Koschaev A.G., Kalyuzhny S.A., Migina E.I., Gavrilenko D.V., Koschaeva O.V. Features of poultry metabolism when using a probiotic feed additive in the diet.
2. Merina-Gluzkina, V.M. (1965). The Comparative Evaluation of Saccharizing vs. Dextrinizing Methods of Determination of Amylase Activity in Intact Humans and Patients with

Acute Pancreatitis. *Russian Clinical Laboratory Diagnostics*, 143.

3. Mikhailova, A.G., Khairullin, R.F., Rumsh, L.D., Demidyuk, I.V., Kostrov, S.V., et al. (2014). Cloning, Sequencing, Expression, and Characterization of Thermostability of Oligopeptidase B from *Serratia proteamaculans*, a Novel Psychrophilic Protease. *Protein Expression and Purification*, 93, 63-76.
4. Batoev, T.Z. (1971). Photometric Analysis of Proteolytic Enzymatic Activities in Pancreas and Its Juice Using the Decrease in Casein Concentration as a Criterion. *The Proceedings of Buryat Agriculture Institute*, 25, 122-126.
5. Vertiprakhov, V.G. and Egorov, I.A. (2016). The Influence of Feed Intake and Conditioned Reflex on Exocrine Pancreatic Function in Broiler Chicks. *Open Journal of Animal Science*, 6, 298-303.
6. Effects of Bile Acids and Lipase Supplementation in Low-Energy Diets on Growth Performance, Fat Digestibility and Meat Quality in Broiler Chickens. *Brazilian Journal of Poultry Science*, 2020, 22 (02), 112-120.
7. Abudabos A. (2014). Effect of fat source, energy level and enzyme supplementation and their interactions on broiler performance. *South African Journal of Animal Science*, 44(3):80-87.
8. WU G. (2018). Principles of animal nutrition. *Boca Raton: CRC Press*.
9. Tancharoenrat P, Ravindran V, Zaefarian F, Ravindran G. (2013). Influence of age on the apparent metabolisable energy and total tract apparent fat digestibility of different fat sources for broiler chickens. *Animal Feed Science and Technology*. 86(3-4):186-192.
10. Siyal FA, Babazadeh D, Wang C, Arain MA, Saeed M, et al. (2017). Emulsifiers in the poultry industry. *World's Poultry Science Journal*. 73(3):611-620.

11. Ravindran V, Tanchaoenrat P, Zaefarian F, Ravindran G. (2016). Fats in poultry nutrition: Digestive physiology and factors influencing their utilisation. *Animal Feed Science and Technology*. 213:1-21.
12. Classen HL. (2017). Diet energy and feed intake in chickens. *Animal Feed Science and Technology*. 233:13-21.
13. Lammasak K, Kijpakorn S, Angkanaporn K. (2019). Corrigendum to: Porcine bile powder supplementation of a high fat broiler diet in relation to growth performance and nutrient digestion. *Animal Production Science* 59(7):1310-1317.
14. Ge XK, Wang AA, Ying ZX, Zhang LG, Su WP, et al. (2018). Effects of diets with different energy and bile acids levels on growth performance and lipid metabolism in broilers. *Poultry Science*. 98(2):887-895.
15. Lai W, Cao A, Li J, Zhang W, Zhang L. (2018). Effect of high dose of bile acids supplementation in broiler feed on growth performance, clinical blood metabolites, and organ development. *The Journal of Applied Poultry Research*. 27(4):532-539.
16. Lai W, Huang W, Dong B, Cao A, Zhang W, et al. (2018). Effects of dietary supplemental bile acids on performance, carcass characteristics, serum lipid metabolites and intestinal enzyme activities of broiler chickens. *Poultry Science*. 97(1):196-202.
17. Nagargoje S, Dhumal MV, Nikam MG, Khose KK. (2016). Effect of crude soy lecithin with or without lipase on performance and carcass traits, meat keeping quality and economics of broiler chicken. *International Journal of Livestock Research*. 6(6):46-54.
18. Hu YD, Lan D, Zhu Y, Pang HZ, Mu XP, (2018). Effect of diets with different energy and lipase levels on performance, digestibility and carcass trait in broilers. *Asian-Australasian Journal of Animal Sciences*. 31(8):1275-1286.
19. Wang Y, Yan J, Zhang X, Han B. (2018). Tolerance properties and growth performance assessment of *Yarrowia lipolytica* lipase in broilers. *Journal of Applied Animal Research*. 46(1):486-491.
20. Meng X, Slominski BA, Guenter W. (2004). The effect of fat type, carbohydrase, and lipase addition on growth performance and nutrient utilization of young broilers fed wheat-based diets. *Poultry Science*. 83(10):1718-1727.
21. R Rej, M Hoder, HU Bergmeyer. (1986). Part. enzymes of concentration catalytic of measurement the for Methods IFCC on recommendation Approved. *Biochem. Clin Chem Clin J*. 32; 510-497.

**Ready to submit your research? Choose ClinicSearch and benefit from:**

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

**At ClinicSearch, research is always in progress.**

Learn more <http://clinicsearchonline.org/journals/international-journal-of-clinical-reports-and-studies>



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.