

Innovation of Audio-Visual Triage system in Combating the Spread of COVID-19 Infection and its efficacy: A Novel Strategy

Muhammad Mansoor Hafeez ^{1*}, Mohammad Azhar ², Hafiz Rizwan Zafar Chudhary ², Muhammad Asim Rana ³, Arif Malik ^{1*}

¹Institute of Molecular Biology and Biotechnology (IMBB), The University of Lahore-Pakistan.

²Accident and Emergency Department, Bahria Town International Hospital, Lahore-Pakistan.

³Intensive Care Unit (ICU), Bahria Town International Hospital, Lahore-Pakistan.

***Corresponding Author:** Muhammad Mansoor Hafeez and Arif Malik. Institute of Molecular Biology and Biotechnology (IMBB), The University of Lahore-Pakistan.

Received Date: July 25, 2023; **Accepted Date:** August 08, 2023; **Published Date:** August 28, 2023

Citation: Muhammad M. Hafeez, Mohammad Azhar, H. R. Zafar Chaudhary, Arif Malik, et al, (2023), Innovation of Audio-Visual Triage system in Combating the Spread of COVID-19 Infection and its efficacy: A Novel Strategy, *J. Clinical Pediatrics and Mother Health*, 2(4); DOI:10.31579/2835-2971/031

Copyright: © 2023, Muhammad Mansoor Hafeez and Arif Malik. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Mus Musculus has provided scientists a well-adapted animal for the developmental studies. There is one on one co-relation between higher mammalian genes to mice genes hence making them superior model to study expression and translation research. Albino strain of house mice were used in this study with comparison of random or blind matings in mice and timed pregnant mice. The working behind blind matings is minimal and maintenance time and required force was also minimal as compared to timed matings. The results of comparisons of weight gain with pregnancies was similar in two groups where some un-pregnant mice also showed average or above average weight gain with time. The accurate embryonic stages requirement for the two experiments were e13.5, e14.5 and e15.5 (peak of neurogenesis in mice). The timed matings showed to be superior regarding total pregnancies and having accurate stages and is recommended to be used in order to save trouble of not getting what you require and more sacrifice of mice then is needed.

Keywords: molecular biology; embryos; human health

Introduction

The pandemics are frightening for everyone, especially for healthcare workers. The most important aspect of dealing with the pandemic is to protect and save the healthcare workers so that they can treat the rest of the people effectively (Mason et al; 2020). Triage is the forefront of an emergency department of an acute hospital which becomes more fundamental during the pandemics. Keeping in view the COVID-19 prevailing situation, a new Triage system to filter and classify the patient's reaching hospital with suspected corona virus infection was developed, named as "AUDIO-VISUAL TRIAGE" system abbreviated as AVT. The recent alarming situation of corona virus infection that has spread to 196 countries around the globe and resulted in failure of routine triage of Emergency Department (Habibi et al; 2020).

Visual Triage System

From the beginning of rapid increase in coronavirus cases every hospital in Pakistan placed a screening triage named as visual triage system outside the main entry door. So, a two-stage triage is introduced, first a visual triage (outside at entry door) & a routine triage (inside Emergency

Department). But at visual triage, healthcare providers felt scared and threatened due to proximity with the patients and their attendants. As a result, they became reluctant and even some of them refused to perform duties at visual triage desk.

Audio-Visual Triage System

Keeping this in mind, an Audio System was added to visual triage that resulted in an AVT. Protocols of AVT included, a medical personal sit on TRIAGE DESK (with glass barrier sheet on it) wearing personal protective equipment (PPE) at a distance of more than 6 feet from the PATIENT DESK. Both desks are connected with Non-touchable MIC SYSTEM for communication (fig 1,2). An Assessment Form with the scoring system is filled by the triage medical staff member and further management of the patient is based on the score achieved (Sivan et al; 2020). In doubtful cases, to reduce the chance of wrong decision (which can be devastating for the whole hospital), an Emergency Doctor re-assess the case and decides accordingly. Then Pathways for the patient management is developed according to the available facilities of the

respective hospital and the instructions of Government. The suspected or confirmed cases are either treated in Isolation wards or referred to respective Centre (designated by The Government) with all the facilities.

Material And Methods

A total of sixty (n=60) participants was recruited to study the effectiveness of AVT System, in the current cross-sectional comparative study. They were further divided into two groups. Participants of group A were performing their duty on visual triage while those of group B were on AVT system for screening the COVID-19 suspected patients. Participants of both groups were provided with the same quality of PPEs. Those participants having previous psychological illness, frequently used to use the washroom, had non-serious attitude and poor compliance with the protocols of wearing PPEs were excluded from the study. After a week every participant was tested by PCR for COVID-19 infection. Questionnaires were distributed among the participants and the response form was collected for further analysis. The study was approved by the ethical review committee of Bahria Town International Hospital Lahore (31.5204° N; 74.3587° E), Pakistan. For the assessment of anxiety, GAD-7 scoring system was used and outcomes were assessed according to the Spitzer et al., 2006 scoring system. The GAD-7 score is calculated by assigning scores of 0, 1, 2, and 3, to the response categories of 'not at all', 'several days', 'more than half the days', and 'nearly every day',

respectively, and adding together the scores for the seven questions. Scores of 5, 10, and 15 are taken as the cut-off points for mild, moderate and severe anxiety, respectively. Using the threshold score of 10, the GAD-7 has a sensitivity of 89% and a specificity of 82% for GAD. It is moderately good at screening three other common anxiety disorders - panic disorder (sensitivity 74%, specificity 81%), social anxiety disorder (sensitivity 72%, specificity 80%) and post-traumatic stress disorder (sensitivity 66%, specificity 81%)

Apart from the GAD-7 scoring, another question, related to job satisfaction, was asked. The data were statistically analyzed with the help of independent t-test using SPSS Software (version 22). In addition, the p-values of less than 0.05 remained statistically significant.

Results

The result of the current study revealed the effectiveness of AVT for the screening of COVID-19 patients. Total of sixty (n=60) participants were allocated equally in two groups. Group A were performing their duties on visual triage with male to female ratio was 21:9 (mean age 28.3 ± 4.01) whereas group B were indulged in screening COVID-19 patients on Audio-Visual Triage (male to female ratio 22:8, mean age=26.56 years). (Table-1). The significantly increased levels of anxiety in group A were found as compared to group B ($p=0.025$).

Variables	Construct	Group a	Group b
Gender	Male	21(70%)	22(73%)
	Female	9(30%)	8(27%)
Age ^a	Minimum	20	20
	Maximum	38	35
Anxiety levels ^b	Anxiety (no)	0(0%)	4(13.3%)
	Anxiety (yes)	30 (100%)	26(86.7%)
	Mild anxiety	6(20%)	15(50%)
	Moderate anxiety	18(60%)	8(26.7%)
	Sever anxiety	6(20%)	3(10%)
Covid-19 pcr ^c	Positive	9(30%)	3(10%)
	Negative	21(70%)	27(90%)
Thinking to quit job	Yes	29(96.7%)	4(13.3%)
	No	1(3.3%)	26(86.7%)

Table 1- Distribution of Demographic and Other Variables Used for The Assessment of The Audio-Visual Triage System

a. Minimum and maximum age in years

b. Anxiety score calculated by GAD-7 scoring system

c. After seven days each participant was test for COVID-19 by PCR

At the end of the study, there were more COVID-19 PCR positive participants in group A ($p=0.001$). (Graph-1). Almost all participants in

group A want to shift their place or ready to quit the job if they were forced to perform the duties at the same visual triage ($p=0.004$). (Table-2)

Variables	¹ group a	² group b	P-value
Anxiety score	^a 3 ± 0.117	2.33 ± 0.154	0.025 ^d
Positive covid-19 pcr	^b 1.30 ± 0.085	1.10 ± 0.056	0.001 ^d
Desire to quit job	^c 1.97 ± 0.033	1.13 ± 0.063	0.004 ^d

Table-2: Outcomes of Different Variables for Assessment of AVT* System

a. Mild Anxiety coded as 1, moderate as 2 and sever as 3

b. COVID-19 Positive PCR coded as 2 and Negative as 1


c. Answer NO coded as 1 and YES as 2

d. independent t-test to find out the significance of variance

1. Thirty participant performing their duties at visual triage screening system

2. Thirty participant performing their duties at AVT system

* AUDIO-VISUAL TRIAGE



Graph-1: Comparison of Triage Systems for COVID-19 Screening

Discussion

Triage is an important area of an Emergency Department of any acute Hospital. In this process, patients are classified according to the severity of their diseases so that the acutely sick and more serious patients are seen before the less serious (Bruijns et al; 2008). Normally, the severity of the disease is assessed in a Triage Room of hospital where the most senior nurse asks about the presenting complaints of patients and checks the vitals of patients before allocating any Category / Class to them. Most of the triage systems have five Categories or Classes. Patients of Category I are the most serious patients, to be seen straight away and patients of Category V are less urgent and would be seen at the last. There are few Triage systems in the world e.g.; Manchester Triage, Australian Triage and Canadian Triage (FitzGerald et al; 2020). But these Triage systems are useful only for classifying patients having routine diseases but cannot be used during disasters like COVID-19 pandemic (Bazyar et al; 2020). If traditional triage criteria are used in pandemics, it can result in the contamination of the Emergency Department and ultimately the whole hospital which may need to be fumigated. Moreover, the Triage staff is liable to be infected and feel threatened which can result in a sensation of frustration and decrease job satisfaction (Forsgårde et al; 2013). An AVT proved to be a simple and effective solution to minimize the elevated psychological stress and more effective triage system in prevention of SARS-CoV-2 infection. Moreover, the pandemics do not give much time for preparation (by the Medical & nonmedical teams) and require big budgets as well as resources by the government. There is a need of increase number of healthcare workers, their training to deal with the pandemic disease, availability of big isolations areas, equipment, accessories and logistics in a short time (Kumar et al; 2020). These become more difficult to manage in poor economic countries. But by taking simple and cost-effective measures like AVT system, for screening suspected subjects, can boost the combating capabilities of frontline healthcare professionals against SARS-CoV-2 virus spread.

Conclusion

The AVT system is more safe and effective strategy to prevent the spread of COVID-19 infection as patients are directed to their respective areas, without compromising the management. So, it can be concluded AVT system should be recommended for all the hospitals dealing with suspected cases of Corona virus infection as it has been found practical, safe and confidence booster for frontline medical fighters on the basis of results of the current study.

Limitations

The residence of participants located at different places like hostels or homes. There would be a chance of virus transmission in participants from their residence.

References

1. Mason, D. J., & Friese, C. R. (2020, March). Protecting health care workers against COVID-19—and being prepared for future pandemics. In *JAMA Health Forum* (Vol. 1, No. 3, pp. e200353-e200353). American Medical Association.
2. Habibi, R., Burci, G. L., de Campos, T. C., Chirwa, D., Cinà, M., Dagron, S., ... & Negri, S. (2020). Do not violate the International Health Regulations during the COVID-19 outbreak. *The Lancet*, 395(10225), 664-666.
3. Sivan, M., Halpin, S., & Gee, J. (2020). Assessing long-term rehabilitation needs in COVID-19 survivors using a telephone screening tool (C19-YRS tool). *Advances in Clinical Neurosciences and Rehabilitation*, 19(4), 14-17.
4. FitzGerald, G., Jelinek, G. A., Scott, D., & Gerditz, M. F. (2010). Emergency department triage revisited. *Emergency Medicine Journal*, 27(2), 86-92.
5. Bruijns, S. R., Wallis, L. A., & Burch, V. C. (2008). A prospective evaluation of the Cape triage score in the emergency department of an urban public hospital in South Africa. *Emergency medicine journal*, 25(7), 398-402.
6. Bazyar, J., Farrokhi, M., & Khankeh, H. (2019). Triage systems in mass casualty incidents and disasters: a review study with a worldwide approach. *Open access Macedonian journal of medical sciences*, 7(3), 482.
7. Kumar, M. S., Raut, R. D., Narwane, V. S., & Narkhede, B. E. (2020). Applications of industry 4.0 to overcome the COVID-19 operational challenges. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 14(5), 1283-1289.
8. Forsgårde, E. S., Attebring, M. F., & Elmqvist, C. (2016). Powerlessness: Dissatisfied patients' and relatives' experience of their emergency department visit. *International emergency nursing*, 25, 32-36.
9. Spitzer, R. L., Kroenke, K., Williams, J. B., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of internal medicine*, 166(10), 1092-1097.

Ready to submit your research? Choose ClinicSearch and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At ClinicSearch, research is always in progress.

Learn more <https://clinicsearchonline.org/journals/clinical-pediatrics-and-mother-health>



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.