

# Awareness and utilization of free maternal healthcare services among women in Mt. Elgon Sub- County, Kenya

Oliver Furechi

IMaseno University School of Arts and Social Sciences, Kisumu, Kenya.

**\*Corresponding Author:** Oliver Furechi, Department of Public Health and Toxicology, Faculty of Veterinary Medicine, University of Nairobi, Kenya.

**Received date:** November 05, 2024; **Accepted date:** November 22, 2024; **Published date:** November 29, 2024

**Citation:** Oliver Furechi, (2024): Awareness and utilization of free maternal healthcare services among women in Mt. Elgon Sub- County, Kenya., *Journal of Heart and Vasculature*, 3(6); **DOI:**10.31579/2834-8788/024

**Copyright:** © 2024, Oliver Furechi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

Maternal mortality remains a matter of public health concern with over 300,000 women dying while giving birth annually. This has driven policy makers to come up with free maternal healthcare services as policy intervention. The Kenyan Government adopted the free maternal healthcare policy in 2013. The policy exempts maternal services from user fees in all public health facilities. This is aimed at promoting skilled delivery to reduce pregnancy-related mortality. In Mt Elgon sub-county, only 45 percent of women deliver in a health facility, this is below the national average of 61.5 percent. This paper presents results from a cross-sectional study conducted in January 2021 in Mt Elgon Sub- County among 377 randomly selected women who delivered in the preceding 12 months and 4 FGDs with women of reproductive age. Descriptive statistics were undertaken while Chi-Square test was used for inferential statistics. Andersen behavioral model of health service use was used to examine how women's demographic characteristics and awareness regarding the free maternal health policy has affected utilization of maternal healthcare services. The results indicate that respondents were generally aware that there is a government policy on free maternal health policy. However, gaps still exist in terms of pushing awareness towards universal reach, infrastructure development, health worker patient ratio, timely provision of essential supplies and long distance to health facilities. This calls for increased sensitization utilizing community health volunteers and allocation of financial and human resources allocation to the department of health within County Governments.

**Keywords:** free health care; maternal; neonatal; cost; Kenya

## Introduction

Maternal mortality remains a matter of public health concern. The world health organization (WHO) estimates that globally 300,000 women die each year with over 90 percent living in low-middle income countries [1].

These preventable mortalities are occasioned by several factors hindering women from accessing and utilizing maternal healthcare services [2] moreover, beyond mortalities, many more women suffer from short and long term disabilities and illness associated with pregnancy such as depression, infertility and fistula [3]. According to WHO, the major causes of maternal deaths comprise; hemorrhage (excessive loss of blood), hypertensive disorders and sepsis/infection which account for more than half of maternal deaths [4]. Most of these deaths could be averted through access to care during pregnancy and delivery period [5]. Access and utilization of antenatal care, skilled birth attendance and postnatal care are therefore critical in reducing maternal mortalities [6,7].

In Kenya, the period 2000 - 2015 saw little or no progress towards achievement of MDG 5 on reducing maternal mortalities. Maternal mortalities remained high at 362 deaths per 100,000 live births, this fell short of the 147/100,000 target [8]. It is against this backdrop that in 2013, the Kenyan Government instituted free maternal health care service to encourage women to take up antenatal, intrapartum, and postnatal healthcare services. The free maternal healthcare service policy is commonly referred to as Linda

mama program, which means protect women. The Linda mama program reimburses health facilities based on their capacity to manage pregnancy and delivery complications. As such, 2500 (\$25) Kenya shillings are reimbursed for every delivery conducted in level 2 facilities (health centers) and level 3 health facilities (sub district hospitals); 5000 (\$50) Kenya shillings are reimbursed for every delivery carried out in level 4 health facilities (district hospitals) and level 5 health facilities (provincial hospitals); and Ksh 17,500 (\$175) is reimbursed for every delivery performed in national referral health facilities [9]. The 2014 Kenya demographic health survey [10] indicates that over 90 percent of women attended at least one antenatal care clinic compared to only 58% that completed the recommended four visits. Moreover, only 62 percent of women deliver with a skilled healthcare provider. Studies [9–12] have identified demographic factors that determine utilization of healthcare service with age, education, socio-economic status, marital status and important bearing in determining utilization of maternal healthcare services [15]. A study in India [16] found that 98% of respondents were aware of different types of family planning methods but only 68% used some family planning method. Furthermore, in Ghana mothers with low knowledge levels on danger signs in pregnancy were less likely to use postnatal care services [17]. While in Uganda utilization of maternal health services varied greatly based on demographic and economic factors of

women with those living far away from towns and with low levels of awareness having low levels of utilization [18]. The studies conducted in Kenya after initiation of free maternal healthcare services focused on implementation of the policy and utilization of the services without considering awareness as a factor that influences utilization [10], [20–22], moreover, within the country disparities in utilization exists [23,24] thus there is need for more nuanced understanding at the local level to inform decisions the newly created County Governments as per the 2010 Constitution. An understanding of such determinants is critical in aiding planning and delivery of maternal healthcare services that is accessible to local communities to improve health outcomes.

## Methods

We conducted a mixed method cross-sectional study that examined how awareness influences utilization of free maternal healthcare services among women. The study was carried out between January and March 2021 in Mt. Elgon Sub County within the larger Bungoma County. The area is part of western Kenya bordering Uganda to the west. It is mountainous and largely inhabited by the Sabaoti ethnic group. The road network is poor with access to some areas using a motorised vehicle virtually impossible. Boda-boda and donkey carts are the common means of transport. The Sub County has 12 health facilities: 10 dispensaries, one sub county hospital and one faith-based health facility. These facilities serve a total population of approximately 65,402. An average of 6200 women delivers in any given year, and 45% of these women deliver within health facilities [10].

The primary study population was 6,327 women who had delivered in the preceding 12 months. Using Yamane (1967) formula a sample of 377 was included in the study. Simple random sampling technique used to identify respondents from a list of 6,327 extracted from community-based health information system (CBHIS). Community health Volunteers (CHVs) guided research assistants to households where selected women stay and did the initial introduction and allowed the research assistants to proceed to administer the semi- structured questionnaire in a private place. Through purposive sampling 5 health managers and 8 CHVs were identified and interviewed as key informants. Additionally, we conducted two FGDs with women who were pregnant or lactating. The semi-structured questionnaire and FGD tools were pretested in a nearby subcounty with similar characteristics to the study area, the tools were revised based on the feedback received.

The independent variable was awareness of free maternal healthcare services while utilization of antenatal service and delivery with skilled birth attendant were the dependent variables. The study was anchored on behavioral model

of health use developed by Andersen [24]. In this framework, individuals' level of awareness is considered as a predisposing factor.

Using Social Package for Social Sciences version 18 quantitative data was analysed to generate descriptive statistics; frequencies, means, percentages while Chi square was used to determine the relationship between independent and dependent variables.  $P < 0.05$  within 95 percent confidence interval was considered significant. Qualitative data was thematically analyzed and reported as verbatim quotes. The study was approved by Maseno University Ethics and Research Committee Approval No. MUERC/00880/20. The Mt. Elgon Sub County health management team allowed the study to be undertaken within their jurisdiction. Study respondents were informed of the objectives of the study and their right to participate voluntarily prior to providing written consent.

## Results

**Socio-demographic and economic characteristics of the respondents** A total of 377 respondents were interviewed. Most of the respondents 208(55.2%) were within the age bracket 26- 34 years, while those within the age group 18-25 were the least represented at 26 (6.9%). In terms of level of education, 251(66.6%) had achieved primary level education with a quarter (24.9%) having attained secondary education, a paltry 4(1.1%) had completed degree level education. Analysis of socio- economic status revealed that more than two-thirds 256(68%) of the respondents belonged to the first and second quintile. A paltry 4.8% (18) were in the fifth quintile see Table 1 for more details. Awareness of free maternal healthcare services

### 1 A motorcycle which is hired for transportation.

A total of 248 (65.8%) were aware of the free maternal health care services. Further analysis revealed that out of the 129 (34.2%) who were not aware, two thirds (68.2%) of them had primary level education, were relatively younger (26-34) and unemployed. A community health volunteer averred. "most of these young girls who get pregnant for the first time are the ones whom we now focus on more to help them understand motherhood and enable them access critical information they need to know".

Furthermore, an analysis based on respondent's social economic status (SES) indicates that all the 18 respondents in the fifth quintile were aware of the existence of free maternal health care services compared to 95 (92.2%) of the respondents within the first quintile. Although the association between SES and awareness was not significant  $\chi^2 (4, N=377) = 7.732 p > .05$ . To support the high levels of awareness, A woman in the FGD said, "We just hear of Linda mama (the name for free maternal health services) from community health volunteers who encourage us to come to health facilities as it is free...we are told it is free"

Variable	Category	Count	%
Age	18-25	26	6.9
	26-34	208	55.2
	Above 35	143	37.9
	Total	377	100.0
Education	Primary	251	66.6
	Secondary	94	24.9
	College	28	7.4
	University	4	1.1
	Total	377	100.0
Marital Status	Married	324	85.9
	Single	46	12.2
	Divorced	7	1.9
	Widowed	0	.0
	Total	377	100.0
SES	First Quintile	103	27.3
	Second Quintile	153	40.6
	Third Quintile	65	17.2

	Fourth Quintile	38	10.1
	Fifth Quintile	18	4.8
	Total	377	100.0

**Table 1: Socio-Demographic and Economic Characteristics of Respondents**

Age of the respondent was not a significant factor determining level of awareness of free maternal health care services  $X^2 (2, N=377) = 3.647, P>.05$ . However, education was significantly associated with awareness  $X^2 (3, N=377) = 8.023, P<.05$ , with all respondents with College or University education being aware of free maternal healthcare services. Moreover, findings indicate that there is no significant association between socioeconomic status with awareness  $X^2 (4, N=377) = 7.732, P>.05$ .

Awareness of specific entitlements provided within the free maternal health care service policy was low. Only half 132 (53.2%) of those who were aware of the free maternal health care services mentioned all the four services, which includes: Antenatal profiling, four antenatal care visit, skilled delivery, and postnatal care. ANC profiling was the least mentioned service entitlement. Even community health volunteers seem not to understand the free maternal healthcare services entitlements very well.

“You know sometimes they (healthcare workers) ask women to pay for the lab...in other cases they don’t...but we usually tell women as part of their preparation to also set aside some money for emergency”. A CHV participant in KII. Moreover, service providers also say the policy is not clear as it does not itemize what it covers, this leads to confusion. One health manager who participated in KII said “A woman can come to the hospital; she is pregnant but on diagnosis she is suffering from non-pregnancy related disease to her she knows she shouldn’t pay because it is free.....we need a clear guidance

and sensitization of community members on what is covered and what is not covered.....some health facilities charge women for ANC profiling (Lab services) while others don’t....it is not clear”. Awareness and uptake of antenatal and skilled birth attendance

Studies [27]– [29] have indicated that there is an association between awareness of healthcare services and utilization. In this study, a cross tabulation of awareness and utilization of four antenatal care services ((ANC4) indicate that those who were aware of free maternal health services were more (65.8%) likely to complete ANC 4 as compared to those who were not aware (34.2%) see table 2 for details. Results of Chi square test of independence indicate that awareness was significantly associated with utilization of four ANC clinics  $X^2 (1, N=377) = 4.85 p<.05$ .

Additionally, attendance of four antenatal care services increases the possibility of delivering in a health facility with a skilled birth attendant. Majority of the respondents 280(74.3%) reported four antenatal care visits while only 27(7.2%) managed to attend one or two clinics during their last pregnancy. A woman who participated in FGD said. “It is important to go even for once so that you can be issued with the booklet {mother baby booklet} so that you are safe and able to use it during delivery and immunization of the baby not to forget the bed net”.

Awareness of Free Maternal Health Care					
		No		Yes	
		Count	%	Count	%
	SBA	72	32.6	149	67.4
Skilled Birth Attendance	Non-SBA	57	36.5	99	63.5
	Total	129	34.2	248	65.8
	Once	8	44.4	10	55.6
	Twice	3	33.3	6	66.7
No. of ANC Clinics attended	Thrice	25	35.7	45	64.3
	Four times	93	33.2	187	66.8
	Total	129	34.2	248	65.8

**Table 2: Awareness and utilization of Antenatal Care and Skilled Birth Attendance services.**

## Discussion

More than 45% of the respondents were not aware of the free maternal healthcare services. This implies that women in this setting are likely to have missed opportunities for critical live saving maternal healthcare services [33]. Unlike the findings Yaya et al. (2017) in their study in Bangladesh, the level of education, socio- economic status was not significantly associated with awareness. Similar findings by are highlighted by Chen et al [30] who concludes that awareness is an important factor that determine demand for preventive medical care among the elderly. Furthermore, Rutaremwa et al., (2015) had a similar finding on education this study differs on their finding on association socioeconomic status with awareness which was not significant. Additionally, Utilization of healthcare services is influenced by the ability of households to cater for the associated costs [34]. In this study, an understanding of household social economic status (SES) unraveled that those in higher socio-economic status have a higher rate of utilization of

antenatal care and skilled delivery as compared to the poor. These results are similar to other studies [19–22] which concluded that despite removal of user fees access to healthcare is highly dependent on individual’s socio-economic status.

The younger cohort in this study (18-25) were fewer among those who completed the recommended four antenatal care visit and delivered with a skilled health provider, these findings reflect the general understanding of young people’s uptake of reproductive health services which is influenced by cognitive and psychosocial barriers. The cognitive barriers are related to awareness of maternal knowledge and available services [36].

Likewise, those above 35 years also had depressed uptake of maternal services, this can be interpreted to imply that women with more than one birth are less likely to use maternal health, this assertion is supported by a reanalysis of demographic health survey data by Mbugua and MacQuarrie 2018 ([37] who found that women above forty years have less odds ratio as

compared to those within 20-29 years bracket. This demonstrates that it is important to focus awareness creation among women who are pregnant for the first time.

Unlike the findings of Gitimu et al., 2015 [38] who concluded that uptake of antenatal care and skilled birth attendance increases as the level of a woman's level of education increases, we found that this situation only applies to those with primary (57.4%) and secondary level (63.8%) education and dropped among those with college/university education (51.8%) in this study.

The implication of low uptake is that women are missing the critical services provided during the four antenatal care visits even health facilities access funds for services not rendered to mothers as the payment is lumpsum and pegged on deliveries reported.

Similar to Ghana's free maternal health policy which was also introduced as a result of poor maternal and neonatal health outcomes and through a political directive that was later operationalized [32], the level of uptake of antenatal care and skilled birth attendance is on the rise [19]. A sustained upward trajectory in uptake of this critical maternal health services will go a long way in accelerating traction towards achievement of SDG 3.

## Conclusion

There was moderate level of women's awareness of free maternal healthcare services that resulted in depressed utilization. Based on these findings, there is need to improve on awareness creation interventions targeting the younger cohorts within the reproductive age. The existing community health volunteers' structures and strengthened facility-based health talks will make a huge difference. Beyond the health system, improving the socio-economic status of the people through programs that target poverty reduction will also have a direct effect on maternal mortalities.

## Acknowledgements

I acknowledge all the respondents who provided valuable insights including health facility managers and community health volunteers. I appreciate Drs. Fred Ikanda and Mary Ochieng' who provided guidance for my Master of Arts in Social Policy at Maseno University.

## References

1. W. H. Organization, Trends in maternal mortality 2000 to 2017: estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division. World Health Organization, 2019.
2. I. A. Ocholla, N. O. Agutu, P. O. Ouma, D. Gatungu, F. O. Makokha, and J. Gitaka, 'Geographical accessibility in assessing bypassing behaviour for inpatient neonatal care, Bungoma County-Kenya', *BMC Pregnancy Childbirth*, vol. 20, no. 1, p. 287, May 2020, doi: 10.1186/s12884-020-02977-x.
3. D. Zelalem Ayele, B. Belayihun, K. Teji, and D. Admassu Ayana, 'Factors Affecting Utilization of Maternal Health Care Services in Kombolcha District, Eastern Hararghe Zone, Oromia Regional State, Eastern Ethiopia', *International Scholarly Research Notices*, Oct. 29, 2014.
4. L. Say et al., 'Global causes of maternal death: a WHO systematic analysis', *Lancet Glob. Health*, vol. 2, no. 6, pp. e323–e333, Jun. 2014.
5. A. Wagstaff and M. Claeson, *The Millennium Development Goals for Health : Rising to the Challenges*. The World Bank, 2004. Accessed: Feb. 03, 2021. [Online].
6. M. Carlough and M. McCall, 'Skilled birth attendance: What does it mean and how can it be measured? A clinical skills assessment of maternal and child health workers in Nepal', *Int. J. Gynecol. Obstet.*, vol. 89, no. 2, pp. 200–208, 2005.
7. M. P. Lambon-Quayefio and N. S. Owoo, 'Examining the Influence of Antenatal Care Visits and Skilled Delivery on Neonatal Deaths in Ghana', *Appl. Health Econ. Health Policy.*, vol. 12, no. 5, pp. 511–522, Oct. 2014, doi: 10.1007/s40258-014-0103-z.
8. E. Lang'at, L. Mwanri, and M. Temmerman, 'Effects of implementing free maternity service policy in Kenya: an interrupted time series analysis', *BMC Health Serv. Res.*, vol. 19, no. 1, p. 645, Sep. 2019.
9. C. M. Gitobu, P. B. Gichangi, and W. O. Mwanda, 'The effect of Kenya's free maternal health care policy on the utilization of health facility delivery services and maternal and neonatal mortality in public health facilities', *BMC Pregnancy Childbirth*, vol. 18, no. 1, Dec. 2018.
10. Kenya National Bureau of Statistics, 'Kenya Demographic and Health Survey 2014', Nairobi, 2014.
11. D. A. Chimankar and H. Sahoo, 'Factors influencing the Utilization of Maternal Health Care Services in Uttarakhand', *Stud. Ethno-Med.*, vol. 5, no. 3, pp. 209–216, Dec. 2011.
12. J. B. Baker and L. Liu, 'The determinants of primary health care utilization: a comparison of three rural clinics in Southern Honduras', *GeoJournal*, vol. 66, no. 4, pp. 295–310, Aug. 2006.
13. D. R. Feikin et al., 'The impact of distance of residence from a peripheral health facility on pediatric health utilisation in rural western Kenya', *Trop. Med. Int. Health*, vol. 14, no. 1, pp. 54–61, 2009, doi: <https://doi.org/10.1111/j.1365-3156.2008.02193.x>.
14. D. Mohan, S. Gupta, A. LeFevre, E. Bazant, J. Killewo, and A. H. Baqui, 'Determinants of postnatal care use at health facilities in rural Tanzania: multilevel analysis of a household survey', *BMC Pregnancy Childbirth*, vol. 15, no. 1, p. 282, Oct. 2015.
15. WHO, 'Barriers and Facilitating Factors in Access to Health Services in the Republic of Moldova', p. 135, 2012.
16. J. R. N. Nansseu, E. C. Nchinda, J.-C. Katte, F. M. Nchagnouot, and G. D. Nguetsa, 'Assessing the knowledge, attitude and practice of family planning among women living in the Mbouda health district, Cameroon', *Reprod. Health*, vol. 12, no. 1, p. 92, Oct. 2015.
17. G. B. Nuamah et al., 'Access and utilization of maternal healthcare in a rural district in the forest belt of Ghana', *BMC Pregnancy Childbirth*, vol. 19, no. 1, p. 6, Dec. 2019.
18. G. Rutaremwa, S. O. Wandera, T. Jhamba, E. Akiror, and A. Kiconco, 'Determinants of maternal health services utilization in Uganda', *BMC Health Serv. Res.*, vol. 15, Jul. 2015.
19. L. M. Calhoun, I. S. Speizer, D. Guilkey, and E. Bukusi, 'The Effect of the Removal of User Fees for Delivery at Public Health Facilities on Institutional Delivery in Urban Kenya', *Matern. Child Health J.*, vol. 22, no. 3, pp. 409–418, 2018.
20. B. B. Masaba and R. M. Mmusi-Phetoe, 'Free Maternal Health Care Policy in Kenya; Level of Utilization and Barriers', *Int. J. Afr. Nurs. Sci.*, vol. 13, p. 100234, Jan. 2020.
21. E. Tama, S. Molyneux, E. Waweru, B. Tsofa, J. Chuma, and E. Barasa, 'Examining the Implementation of the Free Maternity Services Policy in Kenya: A Mixed Methods Process Evaluation', *Int. J. Health Policy Manag.*, vol. 7, no. 7, pp. 603–613, Nov. 2017.
22. T. Achoki et al., 'Health disparities across the counties of Kenya and implications for policy makers, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016', *Lancet Glob. Health*, vol. 7, no. 1, pp. e81–e95, Jan. 2019.
23. J. Mulyanto, A. E. Kunst, and D. S. Kringos, 'Geographical inequalities in healthcare utilisation and the contribution of compositional factors: A multilevel analysis of 497 districts in Indonesia', *Health Place*, vol. 60, p. 102236, Nov. 2019.
24. R. M. Andersen and P. L. Davidson, 'Improving Access to Care in America: Individual and Contextual Indicators.', in *Changing the U.S. health care system: Key issues in health services policy and management*, 3rd ed., San Francisco, CA, US: Jossey-Bass, 2007, pp. 3–31.
25. R. M. Andersen, 'Revisiting the behavioral model and access to medical care: does it matter?', *J. Health Soc. Behav.*, vol. 36, no. 1, pp. 1–10, Mar. 1995.



26. T. Shafqat, S. Fayaz, R. Rahim, and S. Saima, 'Knowledge and awareness regarding antenatal care and delivery among pregnant women', *J. Med. Sci. Peshawar*, vol. 23, pp. 88–91, Apr. 2015.
27. A. K. Ahmed and B. E. Agbana, 'Assessment of awareness, perception and utilization of health services among patients attending clinic at the Muslim Ummah Hospital, Kisi, Southwest, Nigeria', *Niger. J. Med. J. Natl. Assoc. Resid. Dr. Niger.*, vol. 24, no. 1, pp. 37–41, Mar. 2015.
28. S. A. Al-rayes, H. Aldossary, E. Aldoukhi, Z. Alahmedalyousif, G. Aldawood, and A. Alumran, 'The awareness and utilization of 937-telephone health services in Saudi Arabia: Cross-sectional survey study', *Inform. Med. Unlocked*, vol. 20, p. 100393, Jan. 2020.
29. N. Feng, X. Jin, J. Wu, and L. Wang, 'Promotion of awareness and utilization of youth friendly service through multi-sectoral cooperation mechanism in China', *BMC Public Health*, vol. 20, no. 1, p. 747, May 2020.
30. C.-C. Chen, Y.-J. Lin, and Y.-T. Lin, 'Awareness and utilization of preventive care services among the elderly under National Health Insurance', *Int. J. Health Care Finance Econ.*, vol. 13, no. 3–4, pp. 247–260, Dec. 2013.
31. C. Nyongesa, X. Xu, J. J. Hall, W. M. Macharia, F. Yego, and B. Hall, 'Factors influencing choice of skilled birth attendance at ANC: evidence from the Kenya demographic health survey', *BMC Pregnancy Childbirth*, vol. 18, no. 1, p. 88, Apr. 2018.
32. J. Azaare, P. Akweongo, G. C. Aryeetey, and D. Dwomoh, 'Impact of free maternal health care policy on maternal health care utilization and perinatal mortality in Ghana: protocol design for historical cohort study', *Reprod. Health*, vol. 17, no. 1, p. 169, Oct. 2020.
33. E. Åkerman, E. C. Larsson, B. Essén, and R. Westerling, 'A missed opportunity? Lack of knowledge about sexual and reproductive health services among immigrant women in Sweden', *Sex. Reprod. Healthc. Off. J. Swed. Assoc. Midwives*, vol. 19, pp. 64–70, Mar. 2019.
34. M. O. hlarlathie, N. Grede, S. de Pee, and M. Bloem, 'Economic and Social Factors are Some of the Most Common Barriers Preventing Women from Accessing Maternal and Newborn Child Health (MNCH) and Prevention of Mother-to-Child Transmission (PMTCT) Services: A Literature Review', *AIDS Behav.*, vol. 18, no. 5, pp. 516–530, Oct. 2014.
35. F. Masiye, O. Kaonga, and J. M. Kirigia, 'Does User Fee Removal Policy Provide Financial Protection from Catastrophic Health Care Payments? Evidence from Zambia', *PLOS ONE*, vol. 11, no. 1, p. e0146508, Jan. 2016.
36. S. Thongmixay, D. R. Essink, T. de Greeuw, V. Vongxay, V. Sychareun, and J. E. W. Broerse, 'Perceived barriers in accessing sexual and reproductive health services for youth in Lao People's Democratic Republic', *PLOS ONE*, vol. 14, no. 10, p. e0218296, Oct. 2019.
37. S. Mbugua and K. MacQuarrie, 'Determinants of Maternal Care Seeking in Kenya. DHS Further Analysis Reports No. 111.', ICF, 111, 2018. [Online].
38. A. Gitimu et al., 'Determinants of use of skilled birth attendant at delivery in Makueni, Kenya: a cross sectional study', *BMC Pregnancy Childbirth*, vol. 15, no. 1, p. 9, Feb. 2015.
39. T. Asrat, N. Baraki, N. Assefa, and G. Alemkere, 'Birth Preparedness among Women Who Gave Birth in the Last Twelve Months in Jardega Jarte District, Western Ethiopia', *J. Pregnancy*, vol. 2019, pp. 1–8, Apr. 2019.

**Ready to submit your research? Choose ClinicSearch and benefit from:**

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

**At ClinicSearch, research is always in progress.**

Learn more <https://www.clinicsearchonline.org/journals/journal-of-heart-and-vasculature>



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.