

Bempedoic Acid: The Latest in Lipid-lowering

Sanjay Kalra ^{1*}, Navneet Agrawal ²

¹Department of Endocrinology Bharti Hospital, Karnal, Haryana, India

²Department of Diabetology Diabetes Obesity and Thyroid Centre, Gwalior, Madhya Pradesh, India

***Corresponding Author:** Sanjay Kalra, Division of family medicine, Department of Endocrinology Bharti Hospital, Karnal, Haryana, India.

Received date: October 08, 2023; Accepted date: October 23, 2023; Published date: November 08, 2023

Citation: Sanjay Kalra, Navneet Agrawal (2023), Bempedoic Acid: The Latest in Lipid-lowering, *Archives of Clinical and Experimental Pathology*, 2(3); **Doi:**10.31579/2834-8508/019

Copyright: © 2023, Sanjay Kalra. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Bempedoic acid is a novel nonstatin drug that has recently been approved for the management of hypercholesterolemia, in patients with established atherosclerotic cardiovascular disease (ASCVD) or with heterozygous familial hypercholesterolemia (HeFH). This editorial describes the basic and clinical pharmacology of bempedoic acid, and suggests a pragmatic approach to its use in clinical practice.

Dyslipidemia is an important contributor to the burden of atherosclerotic cardiovascular disease (ASCVD). Ensuring optimal lipid levels is an effective way of reducing ASCVD. Conventionally, statins have been used to manage dyslipidemia.¹ However, there still remains a residual risk of cardiovascular disease (CVD) which cannot be mitigated by statins. Some individuals are not able to tolerate the drug, while others do not respond even to maximal doses. Alternatives to statins, such as fibrates, ezetimibe and PCSK9i (proprotein convertase subtilisin/kexin type 9 serine protease inhibitors) are limited by lack of tolerability, efficacy and affordability, respectively.² This is unfortunate, because lipid-lowering is considered low-hanging fruit, for CVD risk reduction, as compared to glucose and weight management. There is a lack of suitable therapies that can serve as complements or alternatives to statins. Keeping this in context, the approval and availability of bempedoic acid in India is a welcome development.

Keywords: bempedoic acid; cholesterol; dyslipidemia; familial hypercholesterolemia; cv risk reduction; residual risk; statin; statin intolerance

1. Interdiction

Bempedoic acid is an orally administered pro-drug with the IUPAC name 8-hydroxy-2,2,14,14-tetramethylpentadecanedioic acid. It inhibits the hepatic enzyme adenosine triphosphate-citrate lyase (ATP-citrate lyase), after being activated to its thioester with coenzyme A, by the enzyme SLC27A2. ATP-citrate lyase acts on hepatic biosynthesis of cholesterol, upstream of the enzyme 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase.³ Thus, its mechanism of action is distinct from, and complementary to, that of statins.

The absorption of bempedoic acid is not affected by food or timing of administration. It reaches peak concentration in 3.5 hours, and has a half-life of 21 ± 11 hours. Excretion is through the renal (70%) and fecal (30%) route.

Clinical Pharmacology

Bempedoic acid has been approved by the United States Food and Drug Administration (USFDA) and the European Medicine Association (EMA). Bempedoic acid has been shown to reduce LDL-C (low-density lipoprotein

cholesterol), total cholesterol, non HDL-C (non-high-density lipoprotein cholesterol) and apolipoprotein B (apo B) levels significantly. In a trial of 2,230 patients with a mean baseline LDL-C of 103.2 ± 29.4 mg/dL, bempedoic acid reduced the mean LDL-C level by 19.2 mg/dL at 12 weeks (-16.5% from baseline).⁴ In the CLEAR (Cholesterol-Lowering via Bempedoic Acid, an ACL-Inhibiting Regimen) Serenity study, statin intolerant patients, with a mean baseline LDL-C of 157.6 mg/dL, bempedoic acid significantly reduced LDL-C (placebo-corrected difference, -21.4%), non-HDL-C (-17.9%), total cholesterol (-14.8%), apo B (-15.0%) and high-sensitivity C-reactive protein (CRP, -24.3% ; $p < 0.001$ for all comparisons).⁵

It can be used as a standalone drug and as part of fixed-dose combination with ezetimibe. It can also be used with statins. It must be noted that concomitant use with bempedoic acid increases blood levels of statins, especially simvastatin and pravastatin. The reason is not known, as bempedoic acid does not interact with the cytochrome P450 enzyme.

Studies are ongoing to determine the effect of bempedoic acid on cardiovascular morbidity and mortality. A systematic review and meta-analysis of 11 trials has shown bempedoic acid to be associated with a reduction in composite cardiovascular outcomes (relative risk [RR] 0.75), along with reduction in LDL-C (21.4%) and CRP (24.3%) levels. In contrast to statins, it is also associated with a reduction in rates of new-onset or worsening of diabetes (RR 0.65).⁶ It is apt to point that bempedoic acid is an AMPK (adenosine monophosphate kinase) activator. Though its clinical significance in the context of lipid-lowering is unknown, AMPK activation is a feature of modern glucose-lowering drugs.

Bempedoic acid inhibits the transporter proteins SLC01B1, SLC01B3 and SLC22A7. Inhibition of SLC22A7, and inhibition of renal tubular OAT2, may lead to a rise in uric acid and precipitate gout. Increase in uric acid usually occurs within the first month of treatment, and persists. The average increase in uric acid at 12 weeks of therapy is 0.8 mg%. The placebo-subtracted risk of gout is 1.1%, and is higher in persons with previous history of gout.⁷ Tendon rupture has been seen in 0.5% patients, and is more common in persons aged >60, those on corticosteroids or fluroquinolones, those with renal failure or previous history of tendon disorders.

Apart from gout, adverse events include muscle spasms, back or limb pain, gout and tendon rupture of the rotator cuff of the shoulder, biceps tendon or Achilles tendon. These adverse effects are rare, however.

Pragmatic Usage

Bempedoic acid can be used in all patients with dyslipidemia, as an adjunct to diet and statins/other lipid-lowering therapy, if existing therapy is unable to achieve the goal. It is also indicated in persons with statin intolerance.³ Bempedoic acid may also be used in heterozygous familial hypercholesterolemia (HeHF), and in persons with ASCVD to lower LDL-C, if it is felt that statin monotherapy will not be able to achieve target LDL-C goal alone.

Table 1 is a list of the clinical indications, concerns, caveats, checkpoints and contraindications associated with bempedoic acid use. The dose is 180 mg orally, once a day, with or without food.

It can be used safely in elderly persons; in those with diabetes irrespective of their glucose-lowering therapy; and with mild-moderate renal or hepatic impairment. It does not have any effect on the QT interval. It can be prescribed with all doses of atorvastatin, rosuvastatin and ezetimibe.

Summary

Bempedoic acid represent a significant breakthrough in lipid management. Current prescribing information and data support its use for dyslipidemia management, in the settings of primary, secondary and tertiary intervention. While initial data is promising, we look forward to real-world evidence in Asian populations, as well as the result of the long-term cardiovascular outcome trial. The availability of bempedoic acid should encourage a change in modern statin-centric lipid management guidelines, and help them evolve into a more person-centered, rather than drug-centered, guidance.

References

1. Michos ED, McEvoy JW, Blumenthal RS. Lipid management for the prevention of atherosclerotic cardiovascular disease. *N Engl J Med*. 2019;381(16):1557-67.
2. Singh M, McEvoy JW, Khan SU, Wood DA, Graham IM, Blumenthal RS, et al. Comparison of transatlantic approaches to lipid management: the AHA/ACC/Multisociety Guidelines vs the ESC/EAS Guidelines. *Mayo Clin Proc*. 2020;95(5):998-1014.
3. Ballantyne CM, Bays H, Catapano AL, Goldberg A, Ray KK, Saseen JJ. Role of bempedoic acid in clinical practice. *Cardiovasc Drugs Ther*. 2021;35(4):853-64.
4. Ray KK, Bays HE, Catapano AL, Lalwani ND, Bloedon LT, Sterling LR, et al; CLEAR Harmony Trial. Safety and efficacy of bempedoic acid to reduce LDL cholesterol. *N Engl J Med*. 2019;380(11):1022-32.
5. Laufs U, Banach M, Mancini GBJ, Gaudet D, Bloedon LT, Sterling LR, et al. Efficacy and safety of bempedoic acid in patients with hypercholesterolemia and statin intolerance. *J Am Heart Assoc*. 2019;8(7):e011662.
6. Wang X, Zhang Y, Tan H, Wang P, Zha X, Chong W, et al. Efficacy and safety of bempedoic acid for prevention of cardiovascular events and diabetes: a systematic review and meta-analysis. *Cardiovasc Diabetol*. 2020;19(1):128.
7. Cicero AFG, Pontremoli R, Fogacci F, Viazzi F, Borghi C. Effect of bempedoic acid on serum uric acid and related outcomes: a systematic review and meta-analysis of the available phase 2 and phase 3 clinical studies. *Drug Saf*. 2020;43(8):727-36.

Ready to submit your research? Choose ClinicSearch and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At ClinicSearch, research is always in progress.

Learn more <https://clinicsearchonline.org/journals/archives-of-clinical-and-experimental-pathology>



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.