

Multivariate Study of Lilium Lancifolium Weed Under Different Field Conditions

Naeem S, Sami A, Haider MZ*, Meeran MW

Department of Plant Breeding and Genetics, Faculty of Agricultural Sciences, University of the Punjab, P.O BOX. 54590, Lahore, Pakistan.

***Corresponding Author:** Haider MZ, Department of Plant Breeding and Genetics, Faculty of Agricultural Sciences, University of the Punjab, P.O BOX. 54590, Lahore, Pakistan.

Received date: October 21, 2024; Accepted date: November 13, 2024; Published date: November 27, 2024

Citation: Naeem S, Sami A, Haider MZ, Meeran MW, (2024), Multivariate Study of Lilium Lancifolium Weed Under Different Field Conditions, *Biomedical and Clinical Research*, 3(6); DOI:10.31579/2834-8486/026

Copyright: © 2024, Haider MZ. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Lilium lancifolium, a weed, is not typically found "around" a specific crop. Still, it is well known for its capacity to adapt to various habitats and compete for scarce resources like water, minerals, and nutrients. To tackle the declining agricultural output caused by this competition, a deliberate study was conducted to examine the characteristics of L. lancifolium in three distinct habitats. The findings revealed a significant relationship between characteristics like height, dry weight, fresh weight, leaf area, leaf length, leaf breadth, and root length. It was interesting to discover that the width of its leaves influenced a plant's height. Additionally, it was shown that location two provided a favorable atmosphere for the ferocious growth and development of L. lancifolium plants. It is highly advised to eradicate or manage L. lancifolium populations to successfully reduce crop output losses brought on by this plant. This study highlights the significance of population management measures.

Keywords: lilium lancifolium; habitats; fresh weight; leaf area; population management

Introduction

Lily (Lilium spp), is a member of the Lilium genus (Huo et al., 2022). The bulbous perennials in the genus Lilium were grown for their enormously beautiful, frequently scented flowers. Due to their stunning beauty, lilies, which have many corolla shapes and colors, were welcomed worldwide. The family Liliaceae includes the genus Lilium, which has about 110–115 species found primarily in East Asia, Europe, and North America in the temperate and frigid parts of the Northern Hemisphere. The various species have four common floral shapes: trumpet, bowl, recurved, and funnel (Kumar et al., 2021). In this study, new populations of a critically endangered Himalayan herb, Lilium polyphyllum D. Don ex Royle, have been reported from Chamba, Himachal Pradesh, India for the first time. Karyotypic, palynological, and serological studies under the light microscope (LM) and scanning electron microscope (SEM) have been carried out on specimens from these populations to add more knowledge to the current species database. Presently studied individuals depicted a diploid (2x) chromosome count of $2n = 24$, confirming the previous Kashmir Himalaya reports. Mitotic study showed a karyotype formula of $2n = 2x = 2m + 2sm + 4st + 16t$, that is, 2 metacentric +2 submetacentric +4 acrocentric +16 telocentric chromosomes. LM and SEM observations revealed that the pollen grains were monosulcate, heteropolar, elliptical in polar view, and oblate with reticulate ornamentation. Tiger lily is a triploid plant with a massive genome that measures 32.8 pg by 47.9 pg (Sun et al., 2021). From spring until October, various lilies bloom, giving

rise to the terms early, mid, and late season (Moore & Lauenroth, 2017). Its diverse habitat includes everything from prairies, meadows, coniferous woods, clearings, and wayside vegetation to coastal scrub. The Columbian lily grows as high as 9,800 feet in lower, drier elevations. The plant blooms for a long time, from early May to early August. The monsoon season and spring, specifically July and August, are the best times for lily blossoming. They need soil that retains moisture and is well-drained. Asiatic, Martagon, Candidum, American, Longiflorum, Trumpet and Aurelian, Oriental, and Other are the 8 categories of hybrids, and there is also a classification for all true species (Kwon et al., 2010). Depending on the type, lilies can be used as cut flowers, in rock gardens, forest gardens, or even containers. Tiger lily, on the other hand, is also regarded as a nutritious food and listed as one of the medicinal plants by the Chinese Ministry of Health (Sun et al., 2022) because it contains a variety of nutrients and bioactive substances, including phenolic glycosides, pectin, steroidal saponins, alkaloids, and different types of vitamins. According to molecular research, lilies first appeared in Eurasia roughly 68 million years ago. Despite their same genetic makeup, Lilium species have developed J. Physical Biomed. Biol. Sci. Volume, 1: 5 Naeem et al., (2022) 2 with noticeable phenotypic variety, which has significant evolutionary relevance. Endlicher (1836) was the first botanist to categorize the entire genus, and he did so by dividing it into five groups based on physical traits: Amblirion, Martagon, Pseudolirium, Eulirion, and Cardiocrinum. The most definitive division of the genus was

suggested by Comber (1949), who divided naturally growing lilies into seven divisions and nine subsections based on 13 morphological traits and two germination types. Martagon Rchb., Pseudolirium Endl., Liriotypus Asch. and Graeb., Archelirion Baker, Sinomartagon Comber, Leucolirion Wilson, and Daurolirion Comber were the seven parts, in that order (Gao et al., 2015). Several techniques have been used to assess the level of genetic diversity, including DNA markers, isozymes, morphological traits, pollen morphology, chromosome traits, and others (Xu et al., 2016). They multiply via seeds and bulbs, which can sprout while still attached to the plant. Digging up the bulbs and carefully collecting all of the clumps is the most efficient technique to eliminate the lilies. All bulbs would need to be dug up, likely 6 to 8 inches below the surface. In Pakistan, farmers employ various techniques to manage the *L. lancifolium* weed. They utilize manual weeding, hoeing, and pulling methods to address small weed infestations, although these approaches demand substantial labor (Khan et al., 2021). Typically, manual practices are combined with other strategies on farms. Conversely, mechanical weed control involves using harrows, cultivators, and rotary tillers to uproot and bury weeds (Bana et al., 2022). While this method is more suitable for larger agricultural operations, it must be executed carefully to prevent soil compaction and crop damage. Chemical weed control is another strategy Pakistani farmers adopt, herbicides like paraquat and 2,4-D specifically for managing *L. lancifolium* {Haider, 2023 #1043}.

Material and Methods

Location of Collection of Samples

The plant samples required for the experimental research were collected from the Faculty of Agricultural Sciences fields.

Collection of Samples

Three locations were allotted for the collection of samples. Ten plants from each site were collected. After taking initial measurements and readings, the plants were carefully put onto a newspaper for drying in sunlight for 2 days. Parameters The parameters studied and recorded are initial plant weight, final plants weight (after drying), plant length, leaf length, leaf area, leaf width, root length, number of inflorescence, initial weight of inflorescence and final weight (after drying). Methods The methods used to measure the parameters including a measuring tape/ruler for length and width and an electronic balance for measuring weight.

Statistics

Statistical analysis is a potent tool for data exploration, testing hypotheses, and model construction in biometry. It offers an organized framework for measuring and analyzing biological systems' patterns, connections, and variations. Researchers can use statistical techniques to identify underlying trends, establish causal linkages, and make predictions based on available data.

ANOVA, correlation analysis, and regression are the statistical techniques used in this study (Nielsen, 2001). We can systematically evaluate data, find patterns, and reach solid findings using ANOVA, correlation, and regression, promoting evidence-based decisionmaking (Cole & Grizzle, 1966).

ANOVA

ANOVA efficiently compares many groups or conditions to spot significant differences and assess the influence of independent variables. (Nielsen, 2001).

Correlation

With the help of correlation analysis, we can determine the degree and direction of an association between two or more variables (Prion & Haerling, 2014).

Regression

Regression analysis enables us to model and predict outcomes based on the relationships between variables, providing a clearer knowledge of the underlying mechanisms at work (McEvoy & Cascio, 1987).

Results and Discussion

The outcomes presented in (Table 1) demonstrate statistically significant differences between the studied locations for *L. lancifolium*. The average dry weight was recorded as $(1.2889 \pm 0.0839\text{cm})$, fresh weight $(5.7422 \pm 0.1081\text{cm})$, number of inflorescence $(21.889 \pm 1.8053\text{g})$, leaf area $(2.0424 \pm 0.5441\text{cm}^2)$, leaf length $(2.9111 \pm 2.911\text{cm})$, leaf width $(0.9222 \pm 0.2244\text{cm})$, plant length $(44.889 \pm 0.8872\text{g})$, and root length $(6.4778 \pm 0.1836\%)$ for *L. lancifolium* plants gathered from the three distinct locations. The plant's fresh weight indicates the resilience of *L. lancifolium* *Lilium lancifolium*, which enables it to withstand various environmental circumstances. The moisture content % further justifies its resistance to hard, hot, and dry weather. Notably, the species' ability to endure dry conditions raises the possibility that it could present crop plants with difficult water, mineral, nutritional, and space constraints, affecting crop yield and productivity. These findings offer insightful information about the competitive and adaptive traits of *L. lancifolium* in various ecological contexts (Li et al., 2023). Tiger lilies, *L. lancifolium*, should be planted in sandy loam soil with a pH range of 6.0 to 7.0 and at least 6 hours in direct sunlight daily for optimum growth (Yang et al., 2021). While their J. Physical Biomed. Biol. Sci. Volume, 1: 5 Naeem et al., (2022) 3 eye-catching and colorful blossoms surely improve a garden's beauty and draw beneficial pollinators like bees and butterflies, it's important to be aware of their potential impact on nearby crops. These negative effects include competing for resources like water and nutrients, shading out smaller plants, having an allelopathic effect that can stunt the growth of neighboring flora, and perhaps luring pests that might spread to nearby crops (Schroth, 1998). Careful planning and administration are essential for successfully overcoming these possible drawbacks. While putting smaller crops in brighter regions can ease the concern about shadowing, strategically spacing the Tiger Lilies and surrounding crops can in decreasing competition for resources. It is possible to promote better coexistence by choosing companion plants resistant to lily allelopathy. The use of appropriate watering methods and mulching strategies not only helps with resource allocation, but also improves the health of the soil (Sahrawat et al., 2010). The risk of pest transmission can be reduced by integrating pest management techniques, such as introducing beneficial insects or applying biological controls. It ensures prompt intervention and upkeep of a peaceful garden to check for illnesses and pests regularly. By taking these steps, gardeners can create a balanced ecosystem where the beauty of Tiger Lilies can be enjoyed alongside thriving neighboring crops.

Table 1. Analysis of variance for morphological traits of *L. lancifolium*

SOV	Fresh weight	Dry weight	Plant Length	Leaf Length	Leaf Width	Leaf Area	Root Length	Inflorescence
Locations	13.0627	0.96444*	706.778	0.80444	0.40111	2.73822	0.30778	56.4444
Error	0.035	0.02111	2.361	0.30778	0.15111	0.88817	0.10111	9.7778
Grand Mean	5.7422	1.2889	44.889	2.9111	0.9222	2.0424	6.4778	21.889
CV	3.26	11.27	3.42	19.06	42.15	46.14	4.91	14.29
Standard Error	0.1081	0.0839	0.8872	0.3203	0.2244	0.5441	0.1836	1.8053

Regression analysis was used to determine the factors that have the greatest influence on plant height. The findings showed that the most important contributing element was leaf width (-0.397332936), which was then followed by leaf area (-0.07707705), dry weight (3.99345008), leaf length (-0.21567442), root length (-0.429935208), inflorescence (-0.244306902), and plant length (0.138632743). Conversely, it was


determined that variables that negatively influenced plant height were moisture % (-0.46), root length (-7.65), and leaf area (-24.20) (Table 3). The following regression equation was developed to predict the relationship: Y (FW) is equal to $-36.606 + 0.003$ (PH), 0.04 (LA), 0.008 (FIW), 9.106 (DW), and 0.424 (MC) {Meeran, 2023 #1044}.

Table 3. Stepwise multiple linear regression for fresh weight of *L. lancifolium*

Traits	Coefficients	Standard Error	t Stat	P-value	Lower 95%	Upper 95%
Leaf width	-0.397332936	1.889940124	-0.210235727	0.868080953	-24.41129909	23.61663322
Leaf Area	-0.077077005	0.910430004	-0.084660001	0.946231982	-11.64518704	11.49103303
Dry weight	3.99345008	0.453782434	8.800362861	0.072031207	-1.772402434	9.759302594
Leaf length	-0.21567442	0.602121238	-0.35819102	0.781032577	-7.866350142	7.435001302
Root length	0.429935208	0.408337386	1.0528921	0.483601326	-4.758483224	5.618353641
Ino	0.244306902	0.068372855	3.573156378	0.173722807	-0.624452588	1.113066392
Plant length	0.138632743	0.027379118	5.063448034	0.124131098	-0.20925194	0.486517426

The Principal of Component Analysis (Figure 1) showed that PC1 predicted only 3% of the variation, while PC2 predicted 95.3 percent of the observed variation in all studied morphological traits, indicating that the majority of the observed differences among the studied traits can be attributed to this single component common factor, which is captured by this component. Additionally, position 3 was found to have the maximum

productivity, which suggests that it is especially conducive to establishing and developing *L. lancifolium* plants. In light of these findings, it is advised to conduct plant population control measures, specifically at position 3. Doing this makes it possible to minimize potential yield losses in agricultural plants and guarantee that *L. lancifolium* grows in the best possible conditions (Clegg et al., 2009).

**Figure 1: Scores Plot for Morphological Traits of *L. lancifolium* across Three Different Locations**

Conclusions

According to the study's findings, site 3 was the most advantageous for the growth and development of *L. lancifolium*. Therefore, it was advised to use methods to limit *L. lancifolium* growth and development to reduce crop output losses.

Data Availability statement All data generated or analyzed during the study have been included in the manuscript.

Ethics approval and consent to participate These aspects are not applicable in this research.

References

- Clegg, S. M., Sklute, E., Dyar, M. D., Barefield, J. E., & Wiens, R. C. (2009). Multivariate analysis of remote laser-induced breakdown spectroscopy spectra using partial least squares, principal component analysis, and related techniques. *Spectrochimica Acta Part B: Atomic Spectroscopy*, 64(1), 79–88. <https://doi.org/10.1016/j.sab.2008.10.045> Cole,

2. J. W. L., & Grizzle, J. E. (1966). Applications of Multivariate Analysis of Variance to Repeated Measurements Experiments. *Biometrics*, 22(4), 810–828. <https://doi.org/10.2307/2528076>
3. J., Zhang, T., Jin, Z.-Y., Xu, X.-M., Wang, J.-H., Zha, X.-Q., & Chen, H.-Q. (2015). Structural characterisation, physicochemical properties and antioxidant activity of polysaccharide from *Lilium lancifolium* Thunb. *Food Chemistry*, 169, 430–438. <https://doi.org/10.1016/j.foodchem.2014.08.016>
- Haider, M.Z., Sami, A., Mazhar, H.S.U.D., Akram, J., NISA, B.U., Umar, M., Meeran, M.W. (2023). Exploring morphological traits variation in *Gomphrena globosa*: A multivariate analysis. *Biol. Agri. Sci. Res.* J., Huo, Y., Li, X., Zhou, J., Lin, S., Ding, J., Liu, H., Yu, J., Hong, N., Wang, G., & Ding, F. (2022). The full-length genome sequence of a novel amalgavirus in *Lilium* spp. In China. *Archives of Virology*, 167(10), 2103–2107. <https://doi.org/10.1007/s00705-022-05523-6>
4. Khan, M. S., Gao, J., Chen, X., Zhang, M., Yang, F., Du, Y., Moe, T. S., Munir, I., Xue, J., & Zhang, X. (2020). Isolation and Characterization of Plant Growth-Promoting Endophytic Bacteria *Paenibacillus polymyxa* SK1 from *Lilium lancifolium*. *BioMed Research International*, 2020, e8650957. <https://doi.org/10.1155/2020/8650957>
5. Meeran, M. W., et al. "Multivariate analysis for morphological traits of *Amaranthus viridis*." *Bulletin of Biological and Allied Sciences Research* 2023.1 (2023): 46-46.
6. Kumar, P., Singh, H., & Kumar Singh, S. (2021). Karyomorphological and SEM studies in newly discovered populations of a critically endangered medicinal plant—*Lilium polyphyllum* D. Don ex Royle from a high altitude protected area of north-western Himalaya. *Microscopy Research and Technique*, 84(10), 2291–2301. <https://doi.org/10.1002/jemt.23784>
7. Kwon, O.-K., Lee, M.-Y., Yuk, J.-E., Oh, S.-R., Chin, Y.-W., Lee, H.-K., & Ahn, K.-S. (2010). Anti- J. Physical Biomed. Biol. Sci. Volume, 1: 5 Naeem et al., (2022) 5 inflammatory effects of methanol extracts of the root of *Lilium lancifolium* on LPS-stimulated Raw264.7 cells. *Journal of Ethnopharmacology*, 130(1), 28–34. <https://doi.org/10.1016/j.jep.2010.04.002>
8. Li, W., Wang, Y., Ren, H., Guo, Z., Li, N., Zhao, C., & Xie, Z. (2023). Transcriptomic and physiological analyses identifying Lanzhou lily (*Lilium davidii* var. *Unicolor*) drought adaptation strategies. *Horticultural Plant Journal*, 9(1), 145–157. <https://doi.org/10.1016/j.hpj.2022.03.003>
9. McEvoy, G. M., & Cascio, W. F. (1987). Do Good or Poor Performers Leave? A Meta-Analysis of the Relationship between Performance and Turnover. *Academy of Management Journal*, 30(4), 744–762. <https://doi.org/10.5465/256158>
10. Moore, L. M., & Lauenroth, W. K. (2017). Differential effects of temperature and precipitation on early- vs. Late-flowering species. *Ecosphere*, 8(5), e01819. <https://doi.org/10.1002/ecs2.1819>
11. Nielsen, R. (2001). Statistical tests of selective neutrality in the age of genomics. *Heredity*, 86(6), Article 6. <https://doi.org/10.1046/j.1365-2540.2001.00895.x>
12. Prion, S., & Haerling, K. A. (2014). Making Sense of Methods and Measurement: Spearman-Rho Ranked-Order Correlation Coefficient. *Clinical Simulation In Nursing*, 10(10), 535–536. <https://doi.org/10.1016/j.ecns.2014.07.005>
13. Sahrawat, K. L., Wani, S. P., Pathak, P., & Rego, T. J. (2010). Managing natural resources of watersheds in the semi-arid tropics for improved soil and water quality: A review. *Agricultural Water Management*, 97(3), 375–381. <https://doi.org/10.1016/j.agwat.2009.10.012>
14. Schroth, G. (1998). A review of belowground interactions in agroforestry, focussing on mechanisms and management options. *Agroforestry Systems*, 43(1), 5–34. <https://doi.org/10.1023/A:1026443018920>
15. Sun, M., Zhao, Y., Shao, X., Ge, J., Tang, X.et.al. (2022). EST–SSR Marker Development and Full-Length Transcriptome Sequence Analysis of Tiger Lily (*Lilium lancifolium* Thunb). *Applied Bionics and Biomechanics*, 2022, e7641048. <https://doi.org/10.1155/2022/7641048>
16. Xu, Z., Wang, H., Wang, B., Fu, L., Yuan, M.et.al. (2016). Characterization and antioxidant activities of polysaccharides from the leaves of *Lilium lancifolium* Thunb. *International Journal of Biological Macromolecules*, 92, 148–155. <https://doi.org/10.1016/j.ijbiomac.2016.07.028>
17. Yang, X., Lu, M., Wang, Y., Wang, Y., Liu, Z., & Chen, S. (2021). Response Mechanism of Plants to Drought Stress. *Horticulturae*, 7(3), Article 3. <https://doi.org/10.3390/horticulturae7030050>

Ready to submit your research? Choose ClinicSearch and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At ClinicSearch, research is always in progress.

Learn more <http://clinicsearchonline.org/journals/biomedical-and-clinical-research>



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.