

Anthropometry of Female Athlete: An Epidemiological Observational Study

MD Shafiullah Prodhania*

Associate Professor, International Institute of Health Sciences.

*Corresponding Author: MD Shafiullah Prodhania, Associate Professor, International Institute of Health Sciences.

Received Date: August 08, 2025; Accepted Date: August 22, 2025; Published Date: September 12, 2025

Citation: MD Shafiullah Prodhania, (2025), Anthropometry of Female Athlete: An Epidemiological Observational Study, *Clinical Trials and Clinical Research*, 4(5); DOI:10.31579/2834-5126/115

Copyright: © 2025, MD Shafiullah Prodhania. This is an open access article distributed under the creative commons' attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Every one of human being is an athlete. In our country many women are serving and exercising regularly. The aim of this research was to explore dietary pattern and nutritional status of female athletes in Bangladesh. This study was a cross-sectional study. About 100 samples were taken randomly and interviewed with pretested structured questionnaire. WHO BMI cut off value for Asian was used. The mean age of respondents was 22 ± 3 years. No underweight and obese was seen. About 76% respondents were normal regarding nutritional status and 24% were overweight. The study result should not be generalized and need further large scale research.

Keywords: anthropometry; athlete; WHO; BMI; RMR

Introduction

Eating habits have influence on athlete's performance. Several factors should be considered in order to plan a suitable nutritional planning, among them the energetic suitability of the diet, the macronutrients distribution and the supply of adequate quantities of vitamins and minerals. Moreover, the athlete's diet should be established according to individual needs, frequency and training intensity and duration [1]. It is known that the high increase of the physical exertion derived from daily exercise and dietetic inadequacy expose the physical activity practitioners to organic problems [2]. Anemia, bone mineral loss and eating disturbs cases related to athletes of both sexes have been registered, as well as amenorrhea, as the main dysfunctions that occur with athletes [3-6]. Generally, eating disturbs are not uniform among athletes, what actually occurs is an expected behavior of deficiencies according to the modality evaluated, especially fights [7], Olympic gymnastics [8] and marathon [9]. Aiming to identify such nutritional deficiencies, one of the proposed strategies is the verification of the energetic consumption and its distribution, besides the macronutrients amount consumed, especially calcium and iron [2]. The aim of this work was to assess anthropometrical measurement of female athletes in Bangladesh.

Methods

This cross-sectional study was conducted at Gazipur and the population of the study was the female athletes of academy. Face to face interview was taken to collect data. Convenient sampling technique was used. SPSS 20.0

version was used to analyze data. Sample was 100 which seem to be representative regarding the study topic. Firstly, a questionnaire was developed in accordance with the study objective to obtain relevant information on the socioeconomic status, food frequency and anthropometric measurement. The questionnaire was pretested. I took interview individually. Our group members asked them the questions and filled the answers and also measured them physically. Weight machine and Height measuring stand were used. After data collection, data were sent to the researcher, which was sorted, scrutinized by the researcher by the selection criteria and then data were analyzed by personal computer by SPSS version 22.0 program.

Results

The mean age of respondents was 22 ± 3 years. Most of them passed SSC examination and married. Mean income of respondents was 7000.44 BDT/month (Table 1). No underweight and obese was seen. About 76% respondents were normal regarding nutritional status and 24% were overweight (Figure 1). In breakfast almost all took ruti/bread, egg, fruits and tea and during lunch rice, fish, dal, leafy vegetables, non-leafy vegetables, potato, meat and fruits were taken 56%, 36%, 62%, 36%, 36%, 20%, 14% and 16% respondents respectively. In dinner rice, dal, vegetable, meat, milk, salad were taken 60%, 50%, 60%, 60%, 56%, 66% respondents respectively (Table 2).

Items	Frequency (%)	Mean (\pm SD)
Mean age in years		22 ± 3

Educational level	
SSC	72(72.0)
HSC	20(20.0)
Graduate	8(8.0)
Marital status	
Single	76(76.0)
Married	24(24.0)
Monthly Income in Taka	7000.44 (± 400)

Table 1: Socio-demographic characteristics of the study subjects.




Figure 1: Anthropometry of respondents.

Food items	Percentage		
	Breakfast	Lunch	Dinner
Rice	0	56	60
Ruti/bread	100	0	20
Fish	0	36	20
Dal	0	62	50
Leafy vegetables	20	36	60
Non leafy vegetables	20	36	60
Potato	0	20	10
Meat	0	14	60
Milk	0	0	56
Egg	100	0	0
Juice	0	0	16
Salad	0	6	66
Fruits	100	16	14
Tea	78	0	0

Table 2: Food habit of respondents.

Discussion

No underweight and obese was seen in present study. About 76% respondents were normal regarding nutritional status and 24% were overweight. But US women, including female athletes, are under ever increasing pressure to be thinner. This pressure to achieve and maintain a low body weight leads to potentially harmful patterns of long-term dieting or disordered eating, which can affect long-term health [10]. Some of the health consequences of long-term energy restriction in female athletes may include poor energy and nutrient intakes, poor nutritional status, decreased RMR and total daily energy expenditure, increased psychological stress and risk for a clinical eating disorder, and increased risk for exercise-induced amenorrhea and osteoporosis [10]. Female athletes participating in thin-build sports may be at risk for the disorders of the female athlete triad: disordered eating, amenorrhea, and osteoporosis [10]. This triad of disorders can also produce severe health consequences that can influence present and future

health [10]. Strategies for helping active women get off the dieting "bandwagon" requires the identification of an appropriate and healthy body weight, good eating and exercise habits, and techniques for maintaining these habits throughout life10. The present study showed that in breakfast almost all took ruti/bread, egg, fruits and tea and during lunch rice, fish, dal, leafy vegetables, non-leafy vegetables, potato, meat and fruits were taken 56%, 36%, 62%, 36%, 36%, 20%, 14% and 16% respondents respectively. In dinner rice, dal, vegetable, meat, milk, salad were taken 60%, 50%, 60%, 60%, 56%, 66% respondents respectively. A study was done to compare the body weight concerns and dieting practices of female collegiate athletes participating in aesthetic, endurance, and team/anaerobic sports. Participants consisted of 425 female athletes from 7 universities across the United States. Body weight concerns and dieting practices were assessed by the Eating Attitudes Test (EAT-26), the Eating Disorder Inventory Body Dissatisfaction Subscale (EDI-BD), and a weight and dieting history questionnaire [11].

Conclusion

Actually, we did study on small sample and this is why study result may not be reflected all over country. To get more precise result large scale study can be done.

Acknowledgment:

The authors express their sincere thanks to all the patients of this study. No external funding was provided for this study.

References

1. (2000). American Dietetic Association, American College Sports Medicine, Dietitians of Canada. Joint Position Statement. Nutrition and athletic performance. *Med Sci Sports Exerc*; 32:2130-2145
2. Cabral CAC, Rosado GP, Silva CHO, Marins JCB. (2006). Diagnosis of the nutritional status of the Weight Lifting Permanent Olympic Team athletes of the Brazilian Olympic Committee (COB). *Rev Bras Med Esporte*;12(6)
3. Hall C, Lane A. (2001). Effects of rapid weight loss on mood and performance among amateur boxers. *Br J Sports Med*; 35:390-395
4. Onywera VO, Kiplamai FK, Boit MK, Pitsiladis YP. (2004). Food and macronutrient intake of elite Kenyan distance runners. *Int J Sport Nutr Exerc Metab*;14(6):709-719
5. Farajian P, Kavouras SA, Yannakoulia M, Sidossis LS. (2004). Dietary intake and nutritional practices of elite Greek aquatic athletes. *Int J Sports Nutr Exerc Metab*; 14(5):574-582
6. Smolak L, Murnen S, Ruble A. (2000). Female athletes and eating problems: a metaanalysis. *Int J Eat Disord*; 27:371-380
7. Umeda T, Nakaji S, Shimoyama T, Yamamoto Y, Totsuka M, et al. (2004). Adverse effects of energy restriction on myogenic enzymes in judoists. *J Sports Sci*; 22:329-338
8. Ribeiro B, Soares E. (2002). Avaliação do estado nutricional de atletas de ginástica olímpica do Rio de Janeiro/ São Paulo. *Revista de Nutrição*; 15:181-91
9. Lehtonen-Veromaa M, Mötönen T, Irljala K, Nuotio I, Leino A, Viikari J. (2000). A 1-year prospective study on the relationship between physical activity, markers of bone metabolism, and bone acquisition in pubertal girls. *J Clin Endocrinol Metab*; 85:3726-32
10. Manore MM. (1999). Nutritional needs of the female athlete. *Clinics in Sports Medicine*; 18(3):549-63
11. Beals KA, Manore MM. (1998). Body Weight Concerns and Dieting Practices of Female Collegiate Athletes. *Journal of The American Dietetic Association - J AMER DIET*; 98(9)

Ready to submit your research? Choose ClinicSearch and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At ClinicSearch, research is always in progress.

Learn more <http://clinicsearchonline.org/journals/clinical-trials-and-clinical-research>